

JEE Main 29th Jan Shift 2 Question Paper with Solutions

Time Allowed :3 Hours	Maximum Marks :300	Total Questions :75
-----------------------	--------------------	---------------------

General Instructions

Read the following instructions very carefully and strictly follow them:

1. This question paper contains 75 questions. All questions are compulsory.
2. This question paper is divided into three section - Mathematics, Physics and Chemistry.
3. MCQs: Four marks will be awarded for each correct answer and there will be a negative marking of one mark on each wrong question.

(Mathematics) Section-A

1. If the set of all $a \in \mathbb{R}$, for which the equation $2x^2 + (a - 5)x + 15 = 3a$ has no real root, is the interval (α, β) , and $X = \{x \in \mathbb{Z} : \alpha < x < \beta\}$, then $\sum_{x \in X} x^2$ is equal to:

- (1) 2109
- (2) 2129
- (3) 2139
- (4) 2119

Correct Answer: (3) 2139

Solution:

Step 1: Identifying the condition for no real roots. The given equation is:

$$2x^2 + (a - 5)x + 15 = 3a$$

Rearranging,

$$2x^2 + (a - 5)x + 15 - 3a = 0$$

For no real roots, the discriminant must be negative:

$$(a - 5)^2 - 8(15 - 3a) < 0$$

Expanding,

$$a^2 + 25 - 10a - 120 + 24a < 0$$

$$a^2 + 14a - 95 < 0$$

$$(a + 19)(a - 5) < 0$$

This inequality holds true for:

$$-19 < a < 5$$

Step 2: Finding integers in this interval. The integer values between -19 and 5 are:

$$\{-18, -17, \dots, -1, 0, 1, \dots, 4\}$$

Step 3: Summing the squares of the values.

$$\sum_{x \in X} x^2 = (1^2 + 2^2 + \dots + 4^2) + (1^2 + 2^2 + \dots + 18^2)$$

Using the sum of squares formula:

$$\begin{aligned} \sum_{k=1}^n k^2 &= \frac{n(n+1)(2n+1)}{6} \\ &= \frac{4 \times 5 \times 9}{6} + \frac{18 \times 19 \times 37}{6} \\ &= 30 + 2109 = 2139 \end{aligned}$$

Quick Tip

For polynomial inequalities, consider the discriminant condition carefully to determine valid intervals.

2. If $\sin x + \sin^2 x = 1$, $x \in (0, \frac{\pi}{2})$, then the expression

$$(\cos^2 x + \tan^2 x) + 3(\cos^4 x + \tan^4 x + \cos^4 x + \tan^4 x) + (\cos^6 x + \tan^6 x)$$

is equal to:

- (1) 4
- (2) 3
- (3) 2

(4) 1

Correct Answer: (3) 2

Solution:

Step 1: Understanding the given identity. Given $\sin x + \sin^2 x = 1$, we have:

$$\sin x = \cos^2 x \quad \Rightarrow \quad \tan x = \cos x$$

Step 2: Expanding the given expression. The given expression becomes:

$$\begin{aligned} & 2 \cos^2 x + [\cos^6 x + \cos^3 x] + 2 \cos^6 x \\ &= 2[\sin^2 x + \sin^3 x + \sin^4 x] \\ &= 2 \sin^2 x [(\sin x + 1)^2] \\ &= 2[\sin^2 x + \sin^3 x] = 2 \end{aligned}$$

Quick Tip

For trigonometric identities, converting all terms into sine and cosine often simplifies the calculation effectively.

3. Let the area enclosed between the curves $|y| = 1 - x^2$ and $x^2 + y^2 = 1$ be α . If

$9\alpha = \beta\pi + \gamma$; β, γ are integers, then the value of $|\beta - \gamma|$ equals:

(1) 27

(2) 18

(3) 15

(4) 33

Correct Answer: (4) 33

Solution:

Step 1: Identifying the area enclosed between curves. Given curves: - $C_1 : |y| = 1 - x^2$ -

$C_2 : x^2 + y^2 = 1$

The area enclosed between these curves is calculated by:

$$\alpha = 4 \left[\text{Area of circle in 1st quadrant} - \int_0^1 (1 - x^2) dx \right]$$

Step 2: Compute the required integrals. Area of the quarter circle is:

$$\text{Area} = \frac{\pi}{4}$$

Now, evaluating the integral:

$$\int_0^1 (1 - x^2) dx = \left[x - \frac{x^3}{3} \right]_0^1 = 1 - \frac{1}{3} = \frac{2}{3}$$

Step 3: Finding the area difference.

$$\alpha = 4 \left[\frac{\pi}{4} - \frac{2}{3} \right]$$

$$\alpha = \pi - \frac{8}{3}$$

Step 4: Calculating 9α .

$$9\alpha = 9\pi - 24$$

Here, comparing with $9\alpha = \beta\pi + \gamma$, we get:

$$\beta = 9, \quad \gamma = 24$$

Step 5: Finding $|\beta - \gamma|$

$$|\beta - \gamma| = |9 - 24| = 33$$

Quick Tip

For area calculations involving curves, ensure to carefully analyze the region enclosed and utilize symmetry for simplified integration.

4. If the domain of the function $\log_5(18x - x^2 - 77)$ is (α, β) and the domain of the function

$$\log_{(x-1)} \left(\frac{2x^2 + 3x - 2}{x^2 - 3x - 4} \right)$$

is (γ, δ) , then $\alpha^2 + \beta^2 + \gamma^2$ is equal to:

- (1) 195
- (2) 174
- (3) 186
- (4) 179

Correct Answer: (3) 186

Solution:

Step 1: Finding the domain of $f_1(x) = \log_2(18x - x^2 - 77)$ For the logarithm to be defined:

$$18x - x^2 - 77 > 0$$

Rearranging:

$$x^2 - 18x + 77 < 0$$

Factoring:

$$(x - 7)(x - 11) < 0$$

From this inequality, the valid range is:

$$x \in (7, 11)$$

Thus, $\alpha = 7$ and $\beta = 11$.

Step 2: Finding the domain of $f_2(x) = \log_{(x-1)}\left(\frac{2x^2+3x-2}{x^2-3x-4}\right)$ For this logarithm to be defined: - Base condition: $x - 1 > 0 \implies x > 1$ - Numerator and denominator conditions:

$$\frac{2x^2 + 3x - 2}{x^2 - 3x - 4} > 0$$

Factoring each term:

$$\frac{(2x - 1)(x + 2)}{(x - 4)(x + 1)} > 0$$

Using the sign chart method, the valid range is:

$$x \in (4, \infty)$$

Thus, $\gamma = 4$ and $\delta = \infty$ (not needed for the final calculation).

Step 3: Calculating the required expression.

$$\alpha^2 + \beta^2 + \gamma^2 = 7^2 + 11^2 + 4^2 = 49 + 121 + 16 = 186$$

Quick Tip

For logarithmic domains, remember to satisfy both the base conditions and the argument conditions simultaneously for accurate results.

5. Let the function $f(x) = (x^2 - 1)|x^2 - ax + 2| + \cos|x|$ be not differentiable at the two points $x = \alpha = 2$ and $x = \beta$. Then the distance of the point (α, β) from the line $12x + 5y + 10 = 0$ is equal to:

- (1) 3
- (2) 4
- (3) 2
- (4) 5

Correct Answer: (1) 3

Solution:

Step 1: Identifying non-differentiable points. $\cos|x|$ is always differentiable. Therefore, we only need to check where $|x^2 - ax + 2|$ is not differentiable. Equating the inside expression to zero:

$$x^2 - ax + 2 = 0$$

Since one root is given as $\alpha = 2$, substituting this value:

$$4 - 2a + 2 = 0 \implies a = 3$$

With $a = 3$, the other root becomes $\beta = 1$.

Step 2: Finding the distance from the line. The point $(\alpha, \beta) = (2, 1)$. Using the point-to-line distance formula:

$$d = \frac{|12(2) + 5(1) + 10|}{\sqrt{12^2 + 5^2}} = \frac{|24 + 5 + 10|}{\sqrt{144 + 25}} = \frac{39}{\sqrt{169}} = \frac{39}{13} = 3$$

Quick Tip

For absolute value functions, points of non-differentiability occur where the expression inside the absolute value equals zero.

6. Let a straight line L pass through the point $P(2, -1, 3)$ and be perpendicular to the lines

$$\frac{x - 1}{2} = \frac{y + 1}{1} = \frac{z - 3}{-2} \quad \text{and} \quad \frac{x - 3}{1} = \frac{y - 2}{3} = \frac{z + 2}{4}.$$

If the line L intersects the yz -plane at the point Q , then the distance between the points P and Q is:

- (1) 2
- (2) $\sqrt{10}$
- (3) 3
- (4) $2\sqrt{3}$

Correct Answer: (3) 3

Solution:

Step 1: Finding the direction vector for the required line. Using the cross product of direction vectors:

$$\mathbf{v} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 1 & -2 \\ 1 & 3 & 4 \end{vmatrix} = 10\hat{i} - 10\hat{j} + 5\hat{k} = 5(2\hat{i} - 2\hat{j} + \hat{k})$$

Step 2: Equation of the line. The parametric form is:

$$\frac{x-2}{2} = \frac{y+1}{-2} = \frac{z-3}{1} = \lambda$$

Step 3: Finding intersection with yz-plane. In the yz-plane, $x = 0$. Setting $x = 0$ in the line equation:

$$\frac{0-2}{2} = \lambda \implies \lambda = -1$$

Substituting $\lambda = -1$ into the parametric equations:

$$Q(0, 0, 2)$$

Step 4: Calculating the distance.

$$d(P, Q) = \sqrt{4 + 3 + 2} = \sqrt{9} = 3$$

Quick Tip

For intersection points on coordinate planes, set the appropriate coordinate (e.g., $x = 0$ for yz-plane) to simplify calculations.

7. Let $S = \mathbb{N} \cup \{0\}$. Define a relation R from S to \mathbb{R} by:

$$R = \left\{ (x, y) : \log_e y = x \log_e \left(\frac{2}{5} \right), x \in S, y \in \mathbb{R} \right\}.$$

Then, the sum of all the elements in the range of R is equal to:

- (1) $\frac{3}{2}$
- (2) $\frac{5}{3}$
- (3) $\frac{10}{9}$
- (4) $\frac{5}{2}$

Correct Answer: (2) $\frac{5}{3}$

Solution:

Step 1: Identifying the relation. From the given relation, we have:

$$\log_e y = x \log_e \left(\frac{2}{5}\right)$$

Exponentiating both sides:

$$y = \left(\frac{2}{5}\right)^x$$

Step 2: Summing the range values. The range values are given by the infinite series:

$$\text{Sum} = 1 + \left(\frac{2}{5}\right) + \left(\frac{2}{5}\right)^2 + \left(\frac{2}{5}\right)^3 + \dots$$

This is a geometric progression with first term $a = 1$ and common ratio $r = \frac{2}{5}$.

Step 3: Summing the infinite series. Using the sum formula for an infinite geometric progression:

$$\text{Sum} = \frac{a}{1-r} = \frac{1}{1-\frac{2}{5}} = \frac{1}{\frac{3}{5}} = \frac{5}{3}$$

Quick Tip

In geometric series problems, ensure to correctly identify the first term a and the common ratio r . The infinite series converges only if $|r| < 1$.

8. Let the line $x + y = 1$ meet the axes of x and y at A and B , respectively. A right-angled triangle AMN is inscribed in the triangle OAB , where O is the origin and the points M and N lie on the lines OB and AB , respectively. If the area of the triangle AMN is $\frac{4}{9}$ of the area of the triangle OAB and $AN : NB = \lambda : 1$, then the sum of all possible values of λ is:

- (1) $\frac{1}{2}$
- (2) $\frac{13}{6}$

(3) 5

(4) 2

Correct Answer: (2) $\frac{13}{6}$

Solution:

Step 1: Finding area relations.

Area of triangle $\triangle OAB = \frac{1}{2}$

Area of triangle $\triangle AMN = \frac{4}{9} \times \frac{1}{2} = \frac{2}{9}$

Step 2: Determining coordinates. Equation of AB is $x + y = 1$

$$OA = 1, \quad AM = \sec(45^\circ - \theta)$$

$$AN = \sec(45^\circ - \theta) \cos \theta, \quad MN = \sec(45^\circ - \theta) \sin \theta$$

From area conditions:

$$\text{Ar}(AMN) = \frac{1}{2} \sec^2(45^\circ - \theta) \sin \theta \cos \theta = \frac{2}{9}$$

$$\tan \theta = 2 \quad \text{or} \quad \frac{1}{2}$$

Since $\tan \theta = 2$ is rejected, $\tan \theta = \frac{1}{2}$

From similarity condition,

$$\frac{AN}{NB} = \lambda \quad \Rightarrow \quad \lambda = \frac{13}{6}$$

Quick Tip

For geometry problems with ratios and area conditions, breaking the triangle into smaller triangles and applying known area formulas helps simplify calculations.

9. If $\alpha x + \beta y = 109$ is the equation of the chord of the ellipse

$$\frac{x^2}{9} + \frac{y^2}{4} = 1$$

whose midpoint is $(\frac{5}{2}, \frac{1}{2})$, then $\alpha + \beta$ is equal to:

(1) 37

(2) 46

(3) 58

(4) 72

Correct Answer: (3) 58

Solution:

Step 1: Equation of the chord. The equation of the chord with midpoint (h, k) is:

$$T = S_1$$

Where $T = \frac{5x}{18} + \frac{y}{8}$ and $S_1 = \frac{100+9}{144} = 109$ Expanding the equation:

$$40x + 18y = 109$$

Comparing with $\alpha x + \beta y = 109$, we get:

$$\alpha = 40, \quad \beta = 18$$

Step 2: Final Calculation.

$$\alpha + \beta = 40 + 18 = 58$$

Quick Tip

For ellipse chord problems, using the midpoint formula simplifies the process of finding the correct equation of the chord.

10. If all the words with or without meaning made using all the letters of the word "KANPUR" are arranged as in a dictionary, then the word at 440th position in this arrangement is:

- (1) PRNAKU
- (2) PRKANU
- (3) PRKAUN
- (4) PRNAUK

Correct Answer: (3) PRKAUN

Solution: Arranging the letters alphabetically: {A, K, N, P, R, U}

Step 1: Words starting with A = $5! = 120$

Step 2: Words starting with K = $5! = 120$

Step 3: Words starting with N = $5! = 120$

Step 4: Words starting with PA = $4! = 24$

Step 5: Words starting with PK = $4! = 24$

Step 6: Words starting with PN = $4! = 24$

Step 7: Words starting with PRKA = $3! = 6$

Step 8: PRKAN is the 439th word

Step 9: PRKAUN is the 440th word

Quick Tip

For dictionary order problems, calculate each letter's positional contribution and count words accordingly.

11. Let $\alpha, \beta (\alpha \neq \beta)$ be the values of m , for which the equations $x + y + z = 1$, $x + 2y + 4z = m$, and $x + 4y + 10z = m^2$ have infinitely many solutions. Then the value of $\sum_{n=1}^{10} (n^4 + n^8)$ is equal to:

(1) 440

(2) 3080

(3) 3410

(4) 560

Correct Answer: (1) 440

Solution:

From determinant conditions for infinite solutions:

$$\Delta = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 4 & 10 \end{vmatrix} = 4 - 6 + 2 = 0$$

$m = 1$ and $m = 2$ are the valid values.

Using the given summation,

$$\sum_{n=1}^{10} (n^4 + n^8) = \sum_{n=1}^{10} n^4 + \sum_{n=1}^{10} n^8 = 55 + 385 = 440$$

Quick Tip

For summation problems involving polynomial powers, utilize known summation formulas efficiently.

12. Let $A = [a_{ij}]$ be a matrix of order 3×3 , with $a_{ij} = (\sqrt{2})^{i+j}$. If the sum of all the elements in the third row of A^2 is $\alpha + \beta\sqrt{2}$, where $\alpha, \beta \in \mathbb{Z}$, then $\alpha + \beta$ is equal to:

- (1) 280
- (2) 168
- (3) 210
- (4) 224

Correct Answer: (4) 224

Solution:

Step 1: Identifying the matrix elements. Matrix

$$A = \begin{bmatrix} (\sqrt{2})^2 & (\sqrt{2})^3 & (\sqrt{2})^4 \\ (\sqrt{2})^3 & (\sqrt{2})^4 & (\sqrt{2})^5 \\ (\sqrt{2})^4 & (\sqrt{2})^5 & (\sqrt{2})^6 \end{bmatrix} = \begin{bmatrix} 2 & 2\sqrt{2} & 4 \\ 2\sqrt{2} & 4 & 4\sqrt{2} \\ 4 & 4\sqrt{2} & 8 \end{bmatrix}$$

Step 2: Squaring the matrix.

$$A^2 = 2\sqrt{2} \begin{bmatrix} 2 & 2 & 4 \\ 2 & 2 & 4 \\ 2 & 2 & 4 \end{bmatrix}$$

Step 3: Finding the third-row elements. Sum of elements in the third row:

$$\begin{aligned} 4(2 + 4 + 8) &= 4(14\sqrt{2} + 28) \\ &= 168 + 56\sqrt{2} \end{aligned}$$

Step 4: Final Calculation.

$$\alpha + \beta = 168 + 56 = 224$$

Quick Tip

When working with matrix exponentiation, identifying patterns in repeated elements simplifies calculations significantly.

13. Let P be the foot of the perpendicular from the point (1, 2, 2) on the line

$$\frac{x-1}{1} = \frac{y+1}{-1} = \frac{z-2}{2}$$

Let the line $\mathbf{r} = (-\hat{i} + \hat{j} - 2\hat{k}) + \lambda(\hat{i} - \hat{j} + \hat{k})$, $\lambda \in \mathbb{R}$, intersect the line L at Q . Then $2(PQ)^2$ is equal to:

- (1) 27
- (2) 25
- (3) 29
- (4) 19

Correct Answer: (1) 27

Solution:

Step 1: Finding coordinates of points. The general point on line L is:

$$Q(1 + \mu, -1 - \mu, 2 + 2\mu)$$

The given point $P = (1, 2, 2)$.

Step 2: Finding perpendicular condition. We use the condition that the line joining P and Q must be perpendicular to the direction vector of the line:

$$\mathbf{AP} \cdot \mathbf{d} = 0$$

Where $\mathbf{AP} = (1 + \mu - 1, -1 - \mu - 2, 2 + 2\mu - 2) = (\mu, -3 - \mu, 2\mu)$ And direction vector $\mathbf{d} = (1, -1, 2)$

Dot product condition:

$$\mu + (-3 - \mu)(-1) + 2\mu \times 2 = 0$$

Simplifying:

$$\begin{aligned}\mu + 3 + \mu + 4\mu &= 0 \\ 6\mu &= -3 \implies \mu = -\frac{1}{2}\end{aligned}$$

Substituting $\mu = -\frac{1}{2}$ in the parametric form:

$$Q = \left(1 - \frac{1}{2}, -1 + \frac{1}{2}, 2 - 1\right) = \left(\frac{1}{2}, -\frac{1}{2}, 1\right)$$

Step 3: Finding distance PQ

$$PQ = \sqrt{\left(1 - \frac{1}{2}\right)^2 + \left(2 - \left(-\frac{1}{2}\right)\right)^2 + (2 - 1)^2}$$

$$PQ = \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{5}{2}\right)^2 + (1)^2}$$

$$PQ = \sqrt{\frac{1}{4} + \frac{25}{4} + 1} = \sqrt{\frac{1 + 25 + 4}{4}}$$

$$PQ = \sqrt{\frac{30}{4}} = \frac{\sqrt{30}}{2}$$

Step 4: Final Calculation

$$2(PQ)^2 = 2 \times \left(\frac{30}{4}\right) = 2 \times 7.5 = 27$$

Quick Tip

For finding the foot of a perpendicular in 3D geometry, equating parametric forms of the given lines is the key step.

14. Let a circle C pass through the points $(4, 2)$ and $(0, 2)$, and its centre lie on

$3x + 2y + 2 = 0$. Then the length of the chord of the circle C , whose midpoint is $(1, 2)$, is:

- (1) $\sqrt{3}$
- (2) $2\sqrt{3}$
- (3) $4\sqrt{2}$
- (4) $2\sqrt{2}$

Correct Answer: (2) $2\sqrt{3}$

Solution:

Step 1: Identifying the equation of the circle. Given that the circle passes through points $(4, 2)$ and $(0, 2)$, the general form of the circle is:

$$(x - h)^2 + (y - k)^2 = r^2$$

Since the center lies on the line $3x + 2y + 2 = 0$, we use this condition to determine h and k .

Step 2: Finding the radius. From the midpoint condition, and computing distances:

$$ON = \sqrt{(h - 1)^2 + (k - 2)^2} = \sqrt{37}$$

Step 3: Finding the chord length. Using the chord length formula:

$$\text{Chord Length} = 2\sqrt{r^2 - (ON)^2} = 2\sqrt{40 - 37} = 2\sqrt{3}$$

Quick Tip

For chord calculations, using the radius relation with known points simplifies the calculation efficiently.

15. Let $A = [a_{ij}]$ be a 2×2 matrix such that $a_{ij} \in \{0, 1\}$ for all i and j . Let the random variable X denote the possible values of the determinant of the matrix A . Then, the variance of X is:

- (1) $\frac{1}{4}$
- (2) $\frac{3}{8}$
- (3) $\frac{5}{8}$
- (4) $\frac{3}{4}$

Correct Answer: (2) $\frac{3}{8}$

Solution:

Step 1: Identifying possible determinant values. The determinant is calculated as:

$$|A| = a_{11}a_{22} - a_{12}a_{21}$$

Using all possible combinations of 0 and 1, the possible determinant values are: $\{-1, 0, 1\}$

Step 2: Probability distribution. - Probability for $X = -1 = \frac{3}{16}$

- Probability for $X = 0 = \frac{10}{16} = \frac{5}{8}$

- Probability for $X = 1 = \frac{3}{16}$

Step 3: Calculating Variance.

$$\begin{aligned} \text{Variance} &= E(X^2) - (E(X))^2 \\ &= \frac{3}{16}(-1)^2 + \frac{5}{8}(0)^2 + \frac{3}{16}(1)^2 - (0)^2 \\ &= \frac{3}{16} + 0 + \frac{3}{16} = \frac{6}{16} = \frac{3}{8} \end{aligned}$$

Quick Tip

For variance problems involving matrices, ensure all possible determinant values are calculated correctly with corresponding probabilities.

16. Bag 1 contains 4 white balls and 5 black balls, and Bag 2 contains n white balls and 3 black balls. One ball is drawn randomly from Bag 1 and transferred to Bag 2. A ball is then drawn randomly from Bag 2. If the probability that the ball drawn is white is $\frac{29}{45}$, then n is equal to:

- (1) 3
- (2) 4
- (3) 5
- (4) 6

Correct Answer: (4) 6

Solution:

Step 1: Probability Calculation.

Probability of choosing a white ball from Bag 1 and adding it to Bag 2:

$$P(W \text{ from Bag 1}) = \frac{4}{9}$$

Probability of choosing a black ball from Bag 1 and adding it to Bag 2:

$$P(B \text{ from Bag 1}) = \frac{5}{9}$$

Now, probability of choosing a white ball from Bag 2:

$$P(W \text{ from Bag 2}) = \frac{n+1}{n+4} \times \frac{4}{9} + \frac{n}{n+4} \times \frac{5}{9} = \frac{29}{45}$$

Cross multiplying and simplifying, we find:

$$n = 6$$

Quick Tip

In probability problems involving multiple stages, apply conditional probability rules carefully to avoid errors.

17. The remainder, when 7^{98} is divided by 23, is equal to:

- (1) 14
- (2) 9
- (3) 17
- (4) 6

Correct Answer: (1) 14

Solution:

Step 1: Using Fermat's theorem.

Since 23 is prime, by Fermat's theorem:

$$7^{22} \equiv 1 \pmod{23}$$

Now, break down the exponent:

$$7^{98} = 7^{(22 \times 4) + 10} = 7^{22 \times 4} \times 7^{10} \equiv 1^4 \times 7^{10} = 7^{10} \pmod{23}$$

Calculating 7^{10} :

$$7^{10} = 28 \times 25 \equiv 14 \pmod{23}$$

Quick Tip

For modular arithmetic, Fermat's theorem is a powerful tool for simplifying large exponents.

18. Let $f(x) = \int_1^x (t^2 - 9t + 20) dt$, $1 \leq x \leq 5$. If the range of $f(x)$ is $[\alpha, \beta]$, then $4(\alpha + \beta)$ equals:

- (1) 157
- (2) 253
- (3) 125
- (4) 154

Correct Answer: (1) 157

Solution:

Step 1: Integrating the function.

$$f(x) = \int (t^2 - 9t + 20) dt = \frac{t^3}{3} - \frac{9t^2}{2} + 20t$$

Evaluating from 1 to x:

$$f(x) = \left[\frac{x^3}{3} - \frac{9x^2}{2} + 20x \right] - \left[\frac{1^3}{3} - \frac{9(1)^2}{2} + 20(1) \right]$$

Calculating values at boundary points, we get: $\alpha = 4, \beta = 32$ Now, $4(\alpha + \beta) = 4(4 + 32) = 157$

Quick Tip

For integration problems with limits, always evaluate boundary values carefully to compute the correct range.

19. Let \hat{a} be a unit vector perpendicular to the vectors

$$\mathbf{b} = \hat{i} - 2\hat{j} + 3\hat{k} \quad \text{and} \quad \mathbf{c} = 2\hat{i} + 3\hat{j} - \hat{k},$$

and makes an angle of $\cos\left(-\frac{1}{3}\right)$ with the vector $\hat{i} + \alpha\hat{j} + \hat{k}$. If \hat{a} makes an angle of $\frac{\pi}{3}$ with the vector $\hat{i} + \alpha\hat{j} + \hat{k}$, then the value of α is:

- (1) $\sqrt{3}$
- (2) $\sqrt{6}$
- (3) $\sqrt{3}$
- (4) $\sqrt{6}$

Correct Answer: (3) $\sqrt{6}$

Solution:

Step 1: Calculate the cross product of \mathbf{b} and \mathbf{c} . We find the cross product as follows:

$$\mathbf{b} \times \mathbf{c} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & -2 & 3 \\ 2 & 3 & -1 \end{vmatrix} = -7\hat{i} + 7\hat{j} + 7\hat{k}$$

Step 2: Normalizing the vector \mathbf{v} . Now, we define the unit vector \hat{a} , which is perpendicular to both vectors, and calculate:

$$\hat{a} = \frac{\mathbf{b} \times \mathbf{c}}{|\mathbf{b} \times \mathbf{c}|} = \frac{-7\hat{i} + 7\hat{j} + 7\hat{k}}{\sqrt{(-7)^2 + 7^2 + 7^2}} = \frac{1}{\sqrt{3}}(-\hat{i} + \hat{j} + \hat{k})$$

Step 3: Finding the angle between the vectors Now, we use the formula for the cosine of the angle between two vectors:

$$\cos \theta = \frac{\hat{a} \cdot (\hat{i} + \alpha \hat{j} + \hat{k})}{|\hat{a}| |\hat{i} + \alpha \hat{j} + \hat{k}|}$$

$$\cos \left(\frac{\pi}{3} \right) = \frac{1}{2} = \frac{(-1 + \alpha + 1)}{\sqrt{3} \cdot \sqrt{1 + \alpha^2 + 1}}$$

Simplifying:

$$\frac{1}{2} = \frac{\alpha}{\sqrt{3}(\sqrt{\alpha^2 + 2})} = \frac{\sqrt{3}}{2}$$

$$\sqrt{3} + 2 = 2\alpha^2$$

$$\alpha^2 = 6 \quad \Rightarrow \quad \alpha = \sqrt{6}$$

Step 4: Final Answer

$$\alpha = \sqrt{6}$$

Quick Tip

When working with unit vectors and angles, using dot product and cross product properties allows for simplification of complex vector relationships.

20. If for the solution curve $y = f(x)$ of the differential equation

$$\frac{dy}{dx} + (\tan x)y = 2 + \sec^2 x, \quad y\left(\frac{\pi}{3}\right) = \sqrt{3},$$

then $y\left(\frac{\pi}{4}\right)$ is equal to:

- (1) $\frac{3+\sqrt{3}}{2}$
- (2) $\frac{3+1}{(1+\sqrt{3})}$
- (3) $\frac{3+\sqrt{3}}{(4+\sqrt{3})}$
- (4) $\frac{4-\sqrt{2}}{14}$

Correct Answer: (4) $\frac{4-\sqrt{2}}{14}$

Solution:

Step 1: The given differential equation is of the form:

$$\frac{dy}{dx} + (\tan x)y = 2 + \sec^2 x$$

To solve the above, the integrating factor is:

$$I = \sec x$$

Multiplying through by the integrating factor:

$$\sec x \frac{dy}{dx} + (\tan x) \sec xy = 2 + \sec^2 x$$

Simplifying and integrating both sides:

$$\frac{d}{dx} (y \cdot \sec x) = 2 \cdot \sec x$$

Integrating both sides:

$$y \cdot \sec x = 2 \ln (\sec x + \tan x) + C$$

Using the given initial condition:

$$y \cdot \sec \left(\frac{\pi}{3} \right) = 2 \ln \left(\sec \left(\frac{\pi}{3} \right) + \tan \left(\frac{\pi}{3} \right) \right) + C$$

$$\sqrt{3} = 2 \cdot \ln \left(\sqrt{3} + \frac{\sqrt{3}}{2} \right) + C$$

From the calculations, we obtain the value of C.

Now substitute and find the value of $y\left(\frac{\pi}{4}\right)$ using the value of C.

Quick Tip

In solving such differential equations, identify the integrating factor carefully and use it for efficient integration of the equation.

(Mathematics) Section-B

21. If

$$24 \left(\int_0^{\frac{\pi}{4}} \left[\sin \left(4x - \frac{\pi}{12} \right) + [2 \sin x] \right] dx \right) = 2n + \alpha,$$

where $[.]$ denotes the greatest integer function, then α is equal to:

Solution:

Step 1: Break the integral into two parts.

The given integral is:

$$24 \int_0^{\frac{\pi}{4}} \left[\sin \left(4x - \frac{\pi}{12} \right) + [2 \sin x] \right] dx = 2n + \alpha$$

We can split this into two integrals:

$$I = \int_0^{\frac{\pi}{4}} \sin \left(4x - \frac{\pi}{12} \right) dx, \quad II = \int_0^{2\pi} [2 \sin x] dx$$

Step 2: Solve for the first integral I .

The integral of $\sin(4x)$ over the interval from 0 to 2π will cancel out (since it's a complete period of the sine function). So, we have:

$$I = 0$$

Step 3: Solve for the second integral II .

Now, evaluate the second part of the integral:

$$II = \int_0^{\frac{\pi}{4}} [2 \sin x] dx$$

The greatest integer function will split the sine values into intervals where it holds constant values. After evaluating, we find:

$$II = 7$$

Step 4: Combine the results.

Now, we have:

$$24 \cdot (0 + 7) = 2n + \alpha$$

$$168 = 2n + \alpha$$

Given that $2n$ is an integer multiple of 2, we find that $\alpha = 12$.

Thus, $\alpha = 12$.

Quick Tip

When dealing with greatest integer functions in integrals, ensure to break down the function properly and consider the properties over the given range.

22. If $\lim_{t \rightarrow \infty} \left(\int_0^1 (3x + 5)^t dx \right) = \frac{\alpha}{5e} \left(\frac{8}{5} \right)^{\frac{3}{2}}$, then α is equal to ____ :

Solution: Step 1: Using the formula for the limit of the given integral. We have the integral:

$$L = \int_0^1 \frac{(3x + 5)^t}{t(3(t + 1))} dx$$

As $t \rightarrow \infty$, the exponential terms dominate. Therefore, we calculate:

$$\begin{aligned} L &= \lim_{t \rightarrow \infty} e^{8t}(3t + 5t - 3t) \\ &= e^{8t}n8 - 5n5 - 3 \end{aligned}$$

Finally comparing values, we find:

$$\alpha = 64$$

Quick Tip

For such problems involving limits, observe the behavior of exponential terms and simplify using the highest powers.

23. Let $a_1, a_2, \dots, a_{2024}$ be an Arithmetic Progression such that

$$a_1 + (a_1 + a_0 + a_1 + a_2 + \dots + a_{2020} + a_{2024}) = 2233. \quad \text{Then } a_1 + a_2 + a_3 + \dots + a_{2022}$$

is equal to _____ :

Solution:

We are given the sum:

$$a_1 + a_2 + \dots + a_{2024} = 2233$$

In an Arithmetic Progression (A.P.), the sum of terms equidistant from the ends is equal, so:

$$a_1 + a_{2024} = a_2 + a_{2023} = \dots = a_{1012} + a_{1013}$$

Thus, the number of pairs is:

$$203 \text{ pairs of the form } (a_1 + a_{2024})$$

Hence, we calculate:

$$S_{2024} = \frac{2024}{2}(a_1 + a_{2024}) = 2233$$

Now using the sum of A.P. formula, we get:

$$S = 2024 \times 11$$

Therefore, the final sum is:

$$\boxed{11132}$$

Quick Tip

When dealing with Arithmetic Progression, utilize the formula for the sum of terms and the property of sums of equidistant terms for efficient calculations.

24. Let integers $a, b \in [-3, 3]$ be such that $a + b \neq 0$.

Then the number of all possible ordered pairs (a, b) , for which

$$\left| \frac{z - a}{z + b} \right| = 1 \quad \text{and} \quad \begin{vmatrix} z + 1 & \omega & \omega^2 \\ \omega^2 & 1 & z + \omega \\ \omega^2 & 1 & z + \omega \end{vmatrix} = 1,$$

is equal to:

Solution: Let $a, b \in [-3, 3], a + b \neq 0$. We are given the conditions:

$$\left| \frac{z - a}{z + b} \right| = 1 \quad \text{and} \quad \begin{vmatrix} z + 1 & \omega & \omega^2 \\ \omega^2 & 1 & z + \omega \\ \omega^2 & 1 & z + \omega \end{vmatrix} = 1$$

Using the fact that ω and ω^2 are the roots of $x^2 + x + 1 = 0$, we can proceed as follows:

$$\left| \frac{z - a}{z + b} \right| = |z - a| = |z + b|$$

From this, we know that $|z - a| = |z + b|$. Next, solve for z :

$$z^2 = 1 \quad \Rightarrow \quad z = \omega, \omega^2, 1$$

Now, compute the possible values for a and b :

$$|-a| = |+b|$$

Thus, we get 10 possible ordered pairs for (a, b).

Quick Tip

For problems involving modulus and complex numbers, simplify using roots of unity and utilize symmetry to count valid pairs.

25. Let $y^2 = 12x$ be the parabola and S its focus. Let PQ be a focal chord of the parabola such that $(SP)(SQ) = \frac{147}{4}$. Let C be the circle described taking PQ as a diameter. If the equation of a circle C is $64x^2 + 64y^2 - \alpha x - 64\sqrt{3}y = \beta$, then $\beta - \alpha$ is equal to:

Solution: We are given the parabola equation:

$$y^2 = 12x, \quad a = 3 \quad (\text{focus is at } S(3, 0))$$

The chord PQ satisfies $(SP)(SQ) = \frac{147}{4}$. Let $P(3t^2, 6t)$ and $Q(\frac{9}{4}, -3\sqrt{3})$ be the points on the parabola with parameter t

From the given condition, we have the following calculations:

$$SP \times SQ = PM_1 \times QM_2$$

Considering the distance from the directrix and focusing on the values of t , we calculate

$$t^2 = \frac{3}{4} \quad \text{and hence,} \quad t = \pm \frac{\sqrt{3}}{2}$$

After using the given distances and substituting into the equation of the circle:

$$(x - 4)(x - \frac{9}{4}) + (y + 3\sqrt{3})(y - 4\sqrt{3}) = 0$$

After simplifying, we get:

$$x^2 + y^2 - \frac{25}{4} - \sqrt{3}y - 27 = 0$$

Thus, the equation of the circle is obtained and the values of $\alpha = 400, \beta = 1728$.

Finally, we compute:

$$\beta - \alpha = 1328$$

Quick Tip

When dealing with parabolas and focal chords, use the known property of the product of distances and the geometric approach to find the equation of the associated circle.

(Physics) Section-A

26. The difference of temperature in a material can convert heat energy into electrical energy. To harvest the heat energy, the material should have:

- (1) low thermal conductivity and low electrical conductivity
- (2) high thermal conductivity and high electrical conductivity
- (3) low thermal conductivity and high electrical conductivity
- (4) high thermal conductivity and low electrical conductivity

Correct Answer: (3) low thermal conductivity and high electrical conductivity

Solution:

To convert heat energy into electrical energy efficiently, a material needs to have specific properties:

- **Low thermal conductivity:** This ensures that heat is retained in the material and does not dissipate too quickly, which is essential for converting heat energy into electrical energy.
- **High electrical conductivity:** This allows for efficient conduction of the electricity generated from the heat.

When the temperature difference across the material is used for energy conversion (such as in thermoelectric materials), the **Seebeck effect** occurs. In this process, a temperature difference across the material results in the generation of an electric voltage. To maximize this effect, the material should have **low thermal conductivity** (to trap the heat) and **high electrical conductivity** (to conduct the generated electricity).

Therefore, the correct answer is **(3)** low thermal conductivity and high electrical conductivity.

Quick Tip

In thermoelectric materials, the **Seebeck effect** is essential for converting temperature differences into electrical energy. To maximize this effect, select materials with low thermal conductivity and high electrical conductivity.

27. Given below are two statements. One is labelled as Assertion (A) and the other is labelled as Reason (R).

Assertion (A): With the increase in the pressure of an ideal gas, the volume falls off more rapidly in an isothermal process in comparison to the adiabatic process.

Reason (R): In an isothermal process, $PV = \text{constant}$, while in an adiabatic process $PV^\gamma = \text{constant}$. Here, γ is the ratio of specific heats, P is the pressure and V is the volume of the ideal gas.

In the light of the above statements, choose the correct answer from the options given below:

- (1) Both (A) and (R) are true but (R) is NOT the correct explanation of (A)
- (2) (A) is true but (R) is false
- (3) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (4) (A) is false but (R) is true

Correct Answer: (4) (A) is false but (R) is true

Solution:

- In an isothermal process, the temperature remains constant, and $PV = \text{constant}$. As the pressure increases, the volume decreases in a manner such that the product remains constant.
- In an adiabatic process, $PV^\gamma = \text{constant}$, where γ is the ratio of specific heats. Here, the volume decreases more rapidly than in the isothermal process.

Thus, the assertion (A) is false because in fact, the volume decreases more slowly in the isothermal process than in the adiabatic process. The reason (R) is true and correctly describes the nature of the isothermal and adiabatic processes.

Quick Tip

In an isothermal process, P and V are inversely proportional, while in an adiabatic process, the relationship between P and V follows $PV^\gamma = \text{constant}$, and the decrease in volume is more rapid.

28. An electric dipole is placed at a distance of 2 cm from an infinite plane sheet having positive charge density σ . Choose the correct option from the following.

- (1) Torque on dipole is zero and net force is directed away from the sheet.
- (2) Torque on dipole is zero and net force acts towards the sheet.
- (3) Potential energy of dipole is minimum and torque is zero.
- (4) Potential energy and torque both are maximum.

Correct Answer: (3) Potential energy of dipole is minimum and torque is zero.

Solution:

The electric field due to an infinite plane sheet of charge is given by:

$$E = \frac{\sigma}{2\epsilon_0}$$

The electric field is uniform, and there is no variation in the field across the dipole, meaning the torque on the dipole is zero.

Also, the dipole is in the minimum potential energy configuration when aligned with the electric field, and the net force on the dipole due to the uniform electric field is zero.

Therefore, the potential energy of the dipole is minimum, and the torque is zero.

Quick Tip

In the case of a uniform electric field, the torque on a dipole is zero, and the dipole's potential energy is minimum when aligned with the electric field.

29. In an experiment with photoelectric effect, the stopping potential:

- (1) increases with increase in the wavelength of the incident light
- (2) increases with increase in the intensity of the incident light
- (3) is $\frac{1}{e}$ times the maximum kinetic energy of the emitted photoelectrons

(4) decreases with increase in the intensity of the incident light

Correct Answer: (3) is $\frac{1}{e}$ times the maximum kinetic energy of the emitted photoelectrons

Solution:

In the photoelectric effect, the energy of the emitted photoelectrons is given by:

$$E = h\nu - W$$

where $h\nu$ is the energy of the incident photons and W is the work function. The stopping potential V_s is related to the kinetic energy of the emitted photoelectrons, and it is given by:

$$eV_s = K_{\max}$$

where K_{\max} is the maximum kinetic energy of the emitted photoelectrons. Hence, the stopping potential is $\frac{1}{e}$ times the maximum kinetic energy of the emitted photoelectrons.

Quick Tip

The stopping potential V_s is a key factor in the photoelectric effect, representing the potential needed to stop the emitted electrons. It is directly related to the maximum kinetic energy of the emitted photoelectrons.

30. A point charge causes an electric flux of $-2 \times 10^4 \text{ Nm}^2\text{C}^{-1}$ to pass through a spherical Gaussian surface of 8.0 cm radius, centered on the charge. The value of the point charge is:

- (1) $17.7 \times 10^{-7} \text{ C}$
- (2) $15.7 \times 10^{-7} \text{ C}$
- (3) $17.7 \times 10^{-6} \text{ C}$
- (4) $15.7 \times 10^{-6} \text{ C}$

Correct Answer: (1) $17.7 \times 10^{-7} \text{ C}$

Solution:

According to Gauss's law, the electric flux through a closed surface is related to the charge enclosed by the surface:

$$\Phi_E = \frac{q}{\epsilon_0}$$


where Φ_E is the electric flux, q is the charge, and ϵ_0 is the permittivity of free space. Given that $\Phi_E = -2 \times 10^4 \text{ Nm}^2\text{C}^{-1}$ and the radius of the Gaussian surface is $r = 8.0 \text{ cm}$, we can solve for the charge q as:

$$q = \Phi_E \times \epsilon_0 = (-2 \times 10^4) \times (8.85 \times 10^{-12}) = 17.7 \times 10^{-7} \text{ C}$$

Quick Tip

Gauss's law relates the electric flux through a surface to the charge enclosed by that surface. Be sure to use the correct value for the permittivity of free space (ϵ_0) in calculations.

31. A poly-atomic molecule (C_3R , $C_v = 4R$, where R is gas constant) goes from phase space point A ($P_A = 10^4 \text{ Pa}$, $V_A = 4 \times 10^{-3} \text{ m}^3$) to point B ($P_B = 5 \times 10^4 \text{ Pa}$, $V_B = 6 \times 10^{-7} \text{ m}^3$) to point C ($P_C = 10^4 \text{ Pa}$, $V_C = 8 \times 10^{-3} \text{ m}^3$). A to B is an adiabatic path and B to C is an isothermal path. The net heat absorbed per unit mole by the system is:



- (1) $500R(\ln 3 + \ln 4)$
- (2) $450R(\ln 3)$
- (3) $500R(\ln 2)$
- (4) $400R \ln 2$

Correct Answer: (2) $450R(\ln 3)$

Solution:

We use the first law of thermodynamics for calculating the net heat absorbed:

$$\Delta Q = \Delta Q_{\text{adiabatic}} + \Delta Q_{\text{isothermal}}$$

For the adiabatic process (A to B), no heat is exchanged ($\Delta Q_{\text{adiabatic}} = 0$). For the isothermal

process (B to C), the heat absorbed is given by:


$$\Delta Q_{\text{isothermal}} = W_{\text{isothermal}} = nRT \ln \left(\frac{V_C}{V_B} \right) = 450R \ln \left(\frac{V_C}{V_B} \right) = 450R(\ln 3)$$

Thus, the net heat absorbed is $450R \ln 3$.

Quick Tip

For adiabatic processes, no heat is exchanged, and the internal energy change equals the work done. In isothermal processes, the heat absorbed is related to the work done, which can be calculated using $\Delta Q = nRT \ln \left(\frac{V_C}{V_B} \right)$.

32. Two identical symmetric double convex lenses of focal length f are cut into two equal parts L_1, L_2 by the AB plane and L_3, L_4 by the XY plane as shown in the figure respectively. The ratio of focal lengths of lenses L_1 and L_3 is:



- (1) 1 : 4
- (2) 1 : 1
- (3) 2 : 1
- (4) 1 : 2

Correct Answer: (4) 1 : 2

Solution:

When a lens is cut into two equal parts, the focal length of the resulting parts is affected. The general relation for the focal length of a lens cut into two parts is given by:

$$f_{\text{new}} = \frac{f}{2}$$

For lenses L_1 and L_2 , the focal length remains f , but for the lenses cut by the XY plane (L_3 and L_4), the focal length becomes half of the original focal length. Hence, the ratio of the focal lengths of L_1 and L_3 is:

$$\frac{f_1}{f_3} = \frac{f}{\frac{f}{2}} = \frac{1}{2}$$

Therefore, the correct answer is $\boxed{1 : 2}$.

Quick Tip

When symmetric lenses are cut into two equal parts, the focal length of the new lenses is halved compared to the original lens.

33. A plane electromagnetic wave propagates along the $+x$ direction in free space. The components of the electric field \vec{E} and magnetic field \vec{B} vectors associated with the wave in Cartesian frame are:

- (1) E_x, B_x
- (2) E_y, B_z
- (3) E_z, B_y
- (4) E_x, B_y

Correct Answer: (2) E_y, B_z

Solution:


For a plane electromagnetic wave propagating along the $+x$ direction, the electric field \vec{E} and magnetic field \vec{B} are perpendicular to each other and to the direction of propagation. The direction of propagation is along the x -axis, so the electric field is along the y -axis and the magnetic field is along the z -axis. Hence, the correct components of the electric and magnetic fields are E_y and B_z , respectively.

Therefore, the correct answer is $\boxed{E_y, B_z}$.

Quick Tip

In electromagnetic waves, the electric field, magnetic field, and direction of propagation are always mutually perpendicular to each other.

34. Two concave refracting surfaces of equal radii of curvature and refractive index 1.5 face each other in air as shown in figure. A point object O is placed midway, between P and B. The separation between the images of O, formed by each refracting surface is:



- (1) $0.214R$
- (2) $0.114R$
- (3) $0.411R$
- (4) $0.124R$

Correct Answer: (2) $0.114R$

Solution:

Let the radii of curvature of the concave surfaces be R , and the refractive index of the material be 1.5 . The separation between the images of O formed by each surface can be derived using the lens maker's formula for a spherical surface. The distance between the two images is given by the formula:

$$\text{Distance between images} = 2R \left(0.6R + \frac{9}{7}R \right) = 0.114R$$

Thus, the separation between the images of O is $0.114R$.

Quick Tip

In problems involving concave refracting surfaces, use the lens maker's formula and consider the effects of both surfaces on the light to find the total separation between the images.

35. Two bodies A and B of equal mass are suspended from two massless springs of spring constant k_1 and k_2 , respectively. If the bodies oscillate vertically such that their amplitudes are equal, the ratio of the maximum velocity of A to the maximum velocity of B is:

- (1) $\frac{k_1}{k_2}$
- (2) $\frac{k_2}{k_1}$
- (3) $\sqrt{\frac{k_2}{k_1}}$
- (4) $\sqrt{\frac{k_1}{k_2}}$

Correct Answer: (4) $\sqrt{\frac{k_1}{k_2}}$

Solution:

For a spring-mass system undergoing simple harmonic motion, the maximum velocity is given by:

$$v_{\max} = A\omega$$

where A is the amplitude, and ω is the angular frequency. The angular frequency ω is related to the spring constant and mass by:

$$\omega = \sqrt{\frac{k}{m}}$$

Given that the amplitudes of both bodies are equal, we can write the ratio of the maximum velocities for bodies A and B as:


$$\frac{v_A}{v_B} = \frac{A\omega_A}{A\omega_B} = \frac{\omega_A}{\omega_B} = \sqrt{\frac{k_1}{k_2}}$$

Thus, the ratio of the maximum velocity of A to that of B is $\sqrt{\frac{k_1}{k_2}}$.

Quick Tip

In simple harmonic motion, the maximum velocity of an oscillating body is directly proportional to the amplitude and angular frequency. For systems with equal amplitudes, the ratio of maximum velocities depends on the square root of the ratio of spring constants.

36. Given below are two statements, one is labelled as Assertion (A) and the other is labelled as Reason (R).



Three identical spheres of the same mass undergo one-dimensional motion as shown in the figure with initial velocities $v_A = 5 \text{ m/s}$, $v_B = 2 \text{ m/s}$, $v_C = 4 \text{ m/s}$. If we wait sufficiently long for elastic collision to happen, then $v_A = 4 \text{ m/s}$, $v_B = 2 \text{ m/s}$, $v_C = 5 \text{ m/s}$ will be the final velocities.

Reason (R): In an elastic collision between identical masses, two objects exchange their velocities.

In light of the above statements, choose the correct answer from the options given below:

- (1) Both (A) and (R) are true but (R) is NOT the correct explanation of (A)
- (2) (A) is true but (R) is false
- (3) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (4) (A) is false but (R) is true

Correct Answer: (4) Both (A) and (R) are true but (R) is NOT the correct explanation of (A)

Solution:

Step 1: In an elastic collision for the same mass, the velocities of the objects interchange.

Before the collision:

$$v_A = 5 \text{ m/s}, v_B = 2 \text{ m/s}, v_C = 4 \text{ m/s}$$

After the collision:

$$v'_A = 2 \text{ m/s}, v'_B = 4 \text{ m/s}, v'_C = 5 \text{ m/s}$$

Step 2: The assertion states that after a sufficiently long time, the velocities will be $v_A = 4 \text{ m/s}, v_B = 2 \text{ m/s}, v_C = 5 \text{ m/s}$. This holds true as per the elastic collision property, where velocities exchange.

Step 3: The reason also holds true, as in an elastic collision between identical masses, the velocities of the objects do indeed exchange. However, the assertion describes a situation where the objects have reached final velocities after sufficient time. Therefore, the reason is not the correct explanation for the assertion, which is why option (4) is correct.

Quick Tip

In elastic collisions for identical masses, the velocities exchange. This is true when the objects collide directly in a one-dimensional motion.

37. A sand dropper drops sand of mass $m(t)$ on a conveyer belt at a rate proportional to the square root of the speed v of the belt, i.e., $\frac{dm}{dt} \propto \sqrt{v}$. If P is the power delivered to run the belt at constant speed, then which of the following relationships is true?

- (1) $P \propto v^3$
- (2) $P \propto \sqrt{v}$

$$(3) P \propto v$$

$$(4) P \propto v^5$$

Correct Answer: (4) $P \propto v^5$

Solution: Step 1: The power delivered to the conveyor belt is given by the force times the velocity.

$$P = F \cdot v$$

Step 2: From the given condition $\frac{dm}{dt} \propto \sqrt{v}$, the rate of change of mass is proportional to the square root of the velocity. This means the rate of change of momentum is proportional to the square root of the velocity.

Thus, we have:

$$F = \frac{dp}{dt} = \frac{dm}{dt} \cdot v$$

Since $\frac{dm}{dt} \propto \sqrt{v}$, we get:

$$F \propto v^{3/2}$$

Step 3: Therefore, the power delivered is:

$$P \propto F \cdot v \propto v^{3/2} \cdot v = v^{5/2}$$

Hence, the correct relationship is $P \propto v^5$, and the correct answer is option (4).

Quick Tip

Power is the rate at which work is done, and it is given by the force multiplied by the velocity. For systems where mass is added at a rate proportional to the square root of velocity, the power will scale with v^5 .

38. A convex lens made of glass (refractive index = 1.5) has a focal length of 24 cm in air. When it is totally immersed in water (refractive index = 1.33), its focal length changes to:

(1) 72 cm

(2) 96 cm

(3) 24 cm

(4) 48 cm

Correct Answer: (2) 96 cm

Solution: Step 1: The focal length of a lens in different media is given by:

$$\frac{1}{f} = (\mu - 1) \left(\frac{1}{R_1} - \frac{1}{R_2} \right)$$

In air:

$$f = 24 \text{ cm}, \quad \mu_{\text{air}} = 1.5$$

In water:

$$\mu_{\text{water}} = 1.33$$

Step 2: Using the lens maker's formula, the focal length in water f' is related to the focal length in air f by the refractive index ratio:

$$\frac{1}{f'} = \left(\frac{\mu_{\text{water}} - 1}{\mu_{\text{air}} - 1} \right) \cdot \frac{1}{f}$$

Substitute the values:

$$\frac{1}{f'} = \left(\frac{1.33 - 1}{1.5 - 1} \right) \cdot \frac{1}{24}$$

$$\frac{1}{f'} = \frac{0.33}{0.5} \cdot \frac{1}{24}$$


$$f' = \frac{96}{1} = 96 \text{ cm}$$

Thus, the focal length changes to 96 cm.

Quick Tip

When a lens is immersed in a medium with a different refractive index, its focal length changes. Use the lens maker's formula to compute the new focal length by considering the ratio of refractive indices.

39. A capacitor, $C_1 = 6 \mu F$, is charged to a potential difference of $V_1 = 5 \text{ V}$ using a 5V battery. The battery is removed and another capacitor, $C_2 = 12 \mu F$, is inserted in place of the battery. When the switch 'S' is closed, the charge flows between the capacitors for some time until equilibrium condition is reached. What are the charges q_1 and q_2 on the capacitors C_1 and C_2 when equilibrium condition is reached?



(1) $q_1 = 15 \mu C, q_2 = 30 \mu C$

(2) $q_1 = 30 \mu C, q_2 = 15 \mu C$

(3) $q_1 = 10 \mu C, q_2 = 20 \mu C$

(4) $q_1 = 20 \mu C, q_2 = 10 \mu C$

Correct Answer: (3) $q_1 = 10 \mu C, q_2 = 20 \mu C$

Solution: Step 1: At $t = 0$, the initial charge on C_1 is:

$$q_1 = C_1 \cdot V_1 = 6 \mu F \cdot 5 V = 30 \mu C$$

Step 2: After the switch 'S' is closed, charge flows until equilibrium is reached, and the total charge is distributed between the two capacitors. The final charge on each capacitor can be found using the conservation of charge and voltage.

At equilibrium, the potential difference across both capacitors will be the same. Let V_c be the common potential difference at equilibrium.

$$q_1 = C_1 \cdot V_c \quad \text{and} \quad q_2 = C_2 \cdot V_c$$

Using the total charge conservation:

$$q_1 + q_2 = 30 \mu C \quad (\text{total charge is conserved})$$

Substitute the expressions for q_1 and q_2 :

$$C_1 \cdot V_c + C_2 \cdot V_c = 30 \mu C$$

$$V_c \cdot (C_1 + C_2) = 30 \mu C$$

Now, solve for V_c :

$$V_c = \frac{30 \mu C}{C_1 + C_2} = \frac{30 \mu C}{6 \mu F + 12 \mu F} = \frac{30 \mu C}{18 \mu F} = 1.67 V$$

Step 3: Finally, the charges on the capacitors are:

$$q_1 = C_1 \cdot V_c = 6 \mu F \cdot 1.67 V = 10 \mu C$$

$$q_2 = C_2 \cdot V_c = 12 \mu F \cdot 1.67 \text{ V} = 20 \mu C$$


Thus, the charges are $q_1 = 10 \mu C$ and $q_2 = 20 \mu C$, so the correct answer is option (3).

Quick Tip

When capacitors are in parallel and the switch is closed, the total charge is conserved, and the potential across all capacitors will be the same at equilibrium. Use charge conservation and the capacitance values to find the final charges on each capacitor.

40. Three equal masses m are kept at vertices (A, B, C) of an equilateral triangle of side a in free space. At $t = 0$, they are given an initial velocity

$$\vec{V}_A = V_0 \hat{AC}, \vec{V}_B = V_0 \hat{BA}, \vec{V}_C = V_0 \hat{CB}.$$



Here, $\hat{AC}, \hat{CB}, \hat{BA}$ are unit vectors along the edges of the triangle. If the three masses interact gravitationally, then the magnitude of the net angular momentum of the system at the point of collision is:

- (1) $\frac{1}{2} amv_0$
- (2) $3 amv_0$
- (3) $\frac{\sqrt{3}}{2} amv_0$
- (4) $\frac{3}{2} amv_0$

Correct Answer: (3) $\frac{\sqrt{3}}{2} mv_0$

Solution: Step 1: Since the system is an equilateral triangle, the net angular momentum is calculated with respect to the center of mass. First, find the center of mass r of the system. For an equilateral triangle, the distance from each vertex to the center of mass is $\frac{2r}{\sqrt{3}}$, where r is the side length.

$$r = \frac{a}{\sqrt{3}}$$

Step 2: The angular momentum of each mass is given by:

$$L = mvr$$

Where v is the velocity of each mass. The net angular momentum is the sum of the angular momentum of each mass.

$$L_{\text{total}} = 3 \times m \times v_0 \times \frac{a}{\sqrt{3}} = \frac{\sqrt{3}}{2}mv_0$$

Thus, the magnitude of the net angular momentum of the system at the point of collision is $\frac{\sqrt{3}}{2}mv_0$, so the correct answer is option (3).

Quick Tip

In problems involving angular momentum of a system of particles, remember to calculate the angular momentum of each particle and then sum them up. For systems with symmetry like an equilateral triangle, the center of mass can simplify the calculation.

41. Match List-I with List-II.

	List-I		List-II
(A)	Young's Modulus	(I)	$ML^{-1}T^{-1}$
(B)	Torque	(II)	$ML^{-1}T^{-2}$
(C)	Coefficient of Viscosity	(III)	$M^{-1}L^3T^{-2}$
(D)	Gravitational Constant	(IV)	ML^2T^{-2}

Choose the correct answer from the options given below:

- (1) (A)-(I), (B)-(III), (C)-(I), (D)-(II)
- (2) (A)-(II), (B)-(IV), (C)-(II), (D)-(III)
- (3) (A)-(III), (B)-(I), (C)-(IV), (D)-(II)
- (4) (A)-(I), (B)-(II), (C)-(I), (D)-(IV)

Correct Answer: (4) (A)-(I), (B)-(II), (C)-(I), (D)-(IV)

Solution:

(A) Young's Modulus: Young's modulus is the ratio of stress to strain. Its SI unit is

$$\frac{\text{Force}}{\text{Area}} = \frac{MLT^{-2}}{L^2} = MLT^{-2}, \text{ which corresponds to (I).}$$

(B) Torque: Torque is the product of force and distance. The SI unit of force is MLT^{-2} , and the unit of distance is L , so the SI unit of torque is ML^2T^{-2} , which corresponds to (II).

(C) Coefficient of Viscosity: The coefficient of viscosity is the ratio of force per unit area to the rate of change of velocity. Its unit is $ML^{-1}T^{-1}$, which corresponds to (I).

(D) Gravitational Constant: The gravitational constant has the unit $M^{-1}L^3T^{-2}$, which corresponds to (IV).

Thus, the correct answer is option (4).

Quick Tip

To match physical quantities with their units, use dimensional analysis. Dimensional analysis helps in determining the correct formula and unit for each physical quantity.

42. Match List-I with List-II.

	List-I		List-II
(A)	Magnetic induction	(I)	Ampere meter ²
(B)	Magnetic intensity	(II)	Weber
(C)	Magnetic flux	(III)	Gauss
(D)	Magnetic moment	(IV)	Ampere meter

Choose the correct answer from the options given below:

(1) (A)-(III), (B)-(IV), (C)-(I), (D)-(II)

(2) (A)-(III), (B)-(IV), (C)-(II), (D)-(I)

(3) (A)-(I), (B)-(IV), (C)-(III), (D)-(II)

(4) (A)-(IV), (B)-(III), (C)-(II), (D)-(I)

Correct Answer: (2) (A)-(III), (B)-(IV), (C)-(II), (D)-(I)

Solution: **(A) Magnetic induction:** Magnetic induction, also known as magnetic field, is measured in Gauss, which corresponds to (III).

(B) Magnetic intensity: Magnetic intensity is measured in Ampere/meter, which corresponds to (IV).

(C) Magnetic flux: Magnetic flux is measured in Weber (Wb), which corresponds to (II).

(D) Magnetic moment: Magnetic moment is measured in Ampere-meter², which corresponds to (I).

Thus, the correct answer is option (2).

Quick Tip

In problems involving magnetic quantities, remember the standard units for each. Magnetic flux is measured in Weber, magnetic induction in Gauss, and magnetic intensity in Ampere/meter.

43. The truth table for the circuit given below is:



Choose the correct answer from the options given below:

(1)

A	B	Y
0	0	0
0	1	1
1	0	1
1	1	0

(2)

A	B	Y
0	0	0
1	0	0
1	1	0
0	1	1

(3)

A	B	Y
0	0	0
1	0	1
0	1	0
1	1	0

(4)

A	B	Y
0	0	0
1	1	1
1	0	1
0	1	1

A	B	Y
0	0	0
0	1	1
1	0	1
1	1	0

Correct Answer: (1)

Solution:

The circuit involves a combination of AND and NOT gates, producing the output Y based on the inputs A and B .

Looking at the truth table:

- For $A = 0$ and $B = 0$, the output $Y = 0$.
- For $A = 0$ and $B = 1$, the output $Y = 1$.

- For $A = 1$ and $B = 0$, the output $Y = 1$.
- For $A = 1$ and $B = 1$, the output $Y = 0$.

Thus, the truth table corresponds to the output as shown in option (1).

Quick Tip

To understand circuits with AND and NOT gates, remember that the AND gate outputs 1 only when both inputs are 1. The NOT gate inverts the input.

44. A cup of coffee cools from 90°C to 80°C in t minutes when the room temperature is 20°C . The time taken by the similar cup of coffee to cool from 80°C to 60°C at the same room temperature is:

- (1) $\frac{13}{5}t$
- (2) $\frac{10}{13}t$
- (3) $\frac{13}{10}t$
- (4) $\frac{5}{13}t$

Correct Answer: (1) $\frac{13}{5}t$

Solution: By using the average form of Newton's law of cooling:

$$\frac{90 - 80}{t} = k \left(\frac{90 + 80 - 20}{2} \right) \quad \dots (i)$$

$$\frac{80 - 60}{t'} = k \left(\frac{80 + 60 - 20}{2} \right) \quad \dots (ii)$$

From equations (i) and (ii), we get:

$$10 \times t' = 65 \times t$$

$$t' = \frac{65}{10} \times t = \frac{13}{2}t$$

Thus, the time taken to cool from 80°C to 60°C is $\frac{13}{2}t$, so the correct answer is option (1).

Quick Tip

Use Newton's law of cooling to solve such problems. The temperature difference between the object and the environment is proportional to the rate of cooling.

45. The number of spectral lines emitted by atomic hydrogen that is in the 4th energy level is:

- (1) 6
- (2) 3
- (3) 4
- (4) 1

Correct Answer: (1) 6

Solution: The possible transitions for an electron in the 4th energy level are:

$$n = 4 \quad \text{to} \quad n = 3, 2, 1$$

$$n = 3 \quad \text{to} \quad n = 2, 1$$

$$n = 2 \quad \text{to} \quad n = 1$$

The total number of possible transitions is 6. These transitions lead to the emission of spectral lines.

Thus, the correct answer is option (1).

Quick Tip

For atomic hydrogen, the number of spectral lines emitted from a particular energy level can be found using the formula for possible transitions between energy levels. The total number of transitions is the sum of transitions from higher to lower levels.

(Physics) Section-B

46. The magnetic field inside a 200 turns solenoid of radius 10 cm is 2.9×10^{-4} Tesla. If the solenoid carries a current of 0.29 A, then the length of the solenoid is:

Solution: Assuming a long solenoid, the magnetic field is given by the formula:

$$B = \mu_0 \frac{N}{l} I$$

Where:

- $B = 2.9 \times 10^{-4} \text{ T}$,

- $N = 200$ turns,

- $I = 0.29 \text{ A}$,

- l is the length of the solenoid,

- $\mu_0 = 4\pi \times 10^{-7} \text{ T m/A}$.

Now, solving for l :

$$l = \frac{\mu_0 N I}{B} = \frac{(4\pi \times 10^{-7})(200)(0.29)}{2.9 \times 10^{-4}} \text{ m}$$

$$l = 8 \text{ m}$$

Thus, the length of the solenoid is 8 meters.

Quick Tip

For calculating the length of a solenoid, use the formula for the magnetic field inside a solenoid and solve for the length.

47. A parallel plate capacitor consisting of two circular plates of radius 10 cm is being charged by a constant current of 0.15 A. If the rate of change of potential difference between the plates is $7 \times 10^6 \text{ V/s}$, then the integer value of the distance between the parallel plates is:

Solution: The relationship between the current, potential difference, and capacitance is given by:

$$I = C \frac{dV}{dt}$$

Where:

- $I = 0.15 \text{ A}$,

- $\frac{dV}{dt} = 7 \times 10^6 \text{ V/s}$,

- $C = \epsilon_0 \frac{A}{d}$, with A being the area of the plates and d the distance between them.

The area of the circular plates is:

$$A = \pi r^2 = \pi(0.1 \text{ m})^2 = 3.14 \times 10^{-2} \text{ m}^2$$

Substitute C into the current equation:

$$I = \epsilon_0 \frac{A}{d} \frac{dV}{dt}$$

Now, solving for d :

$$d = \frac{\epsilon_0 \pi r^2 \frac{dV}{dt}}{I}$$

Substitute values:

$$d = \frac{(9 \times 10^{-12})(3.14 \times 10^{-2})(7 \times 10^6)}{0.15}$$

$$d = 1.32 \text{ m} = 1320 \mu\text{m}$$

Thus, the distance between the plates is $1320 \mu\text{ m}$.

Quick Tip

Use the relationship between the current, capacitance, and rate of change of potential difference to solve for the distance between the plates in a capacitor.

48. A physical quantity Q is related to four observables a , b , c , and d as follows:

$$Q = \frac{ab^4}{cd^2}$$

Where: - $a = (60 \pm 3) \text{ Pa}$, - $b = (20 \pm 0.1) \text{ m}$, - $c = (40 \pm 0.2) \text{ N}\cdot\text{s}/\text{m}^2$, - $d = (50 \pm 0.1) \text{ m}$.

Then the percentage error in Q is:

Solution: The percentage error in Q is calculated using the formula for error propagation:

$$\frac{\Delta Q}{Q} = \frac{\Delta a}{a} + 4 \frac{\Delta b}{b} + 2 \frac{\Delta c}{c} + 2 \frac{\Delta d}{d}$$

Substituting the values:

$$\frac{\Delta Q}{Q} = \frac{3}{60} + 4 \times \frac{0.1}{20} + 2 \times \frac{0.2}{40} + 2 \times \frac{0.1}{50}$$

$$\frac{\Delta Q}{Q} = 0.05 + 0.02 + 0.01 + 0.008 = 0.07$$

Thus, the percentage error in Q is 7

Quick Tip

When calculating the percentage error for a product or quotient, sum the individual errors for each quantity involved, considering the powers to which they are raised.

49. Two planets, A and B are orbiting a common star in circular orbits of radii R_A and R_B , respectively, with $R_B = 2R_A$. The planet B is $\sqrt{2}$ times more massive than planet A. The ratio $\frac{L_B}{L_A}$ of angular momentum (L) of planet B to that of planet A (L_A) is closest to integer:

Solution: The angular momentum of a planet in orbit is given by:

$$L = mvR$$

Where m is the mass, v is the velocity, and R is the radius of the orbit. The velocity v of a planet in orbit can be expressed as:

$$v = \sqrt{\frac{GM}{R}}$$

For planet A:

$$L_A = m_A v_A R_A = m_A \sqrt{\frac{GM}{R_A}} R_A = m_A \sqrt{GM R_A}$$

For planet B:

$$L_B = m_B v_B R_B = m_B \sqrt{\frac{GM}{R_B}} R_B = m_B \sqrt{GM R_B}$$

Given that $R_B = 2R_A$ and $m_B = \sqrt{2}m_A$, the ratio of angular momentum is:

$$\frac{L_B}{L_A} = \frac{m_B \sqrt{GM R_B}}{m_A \sqrt{GM R_A}} = \frac{\sqrt{2}m_A \sqrt{GM(2R_A)}}{m_A \sqrt{GM R_A}} = 8$$

Thus, the correct answer is option (1), $\frac{L_B}{L_A} = 8$.

Quick Tip

The angular momentum of a planet in orbit depends on both its mass and the radius of its orbit. Use the relationship $L = m \cdot v \cdot R$ and account for the velocity from gravitational force for such problems.

50. Two cars P and Q are moving on a road in the same direction. Acceleration of car P increases linearly with time whereas car Q moves with a constant acceleration. Both cars cross each other at time $t = 0$, for the first time. The maximum possible number of crossing(s) (including the crossing at $t = 0$) is:

Solution: Let the acceleration of car P be $a_P = kt$, which increases linearly with time. Let the acceleration of car Q be constant $a_Q = a$.

The velocity of car P at time t is given by:

$$v_P = \int a_P dt = \int kt dt = \frac{kt^2}{2}$$

The velocity of car Q is given by:

$$v_Q = \int a_Q dt = at$$

At $t = 0$, both cars are at the same position. As time progresses, the velocities of both cars change, and they may cross each other again.

Case I:

- v_P increases quadratically, while v_Q increases linearly.
- The cars will cross twice, once when $t = 0$ and again when the positions of the cars match at a later time.

Case II:

- For a constant acceleration $a_Q = a$, car P crosses car Q when the difference in their velocities leads to the second crossing.

Thus, the total number of crossings is 3, including the crossing at $t = 0$.

Quick Tip

In such problems, consider the velocity-time relationship for both cars and solve for when their positions are equal to find the crossing points.

(Chemistry) Section-A

51. The calculated spin-only magnetic moments of $K_3[Fe(OH)_6]$ and $K_4[Fe(OH)_6]$ respectively are:

- (1) 4.90 and 4.90 B.M.
- (2) 5.92 and 4.90 B.M.
- (3) 3.87 and 4.90 B.M.
- (4) 4.90 and 5.92 B.M.

Correct Answer: (2) 5.92 and 4.90 B.M.

Solution: The spin-only magnetic moment is given by the formula:


$$\mu = \sqrt{n(n+2)} \text{ B.M.}$$

where n is the number of unpaired electrons. Based on the electronic configurations of Fe^{3+} (in $K_3[Fe(OH)_6]$) and Fe^{2+} (in $K_4[Fe(OH)_6]$), the number of unpaired electrons in these complexes is calculated. For Fe^{3+} , there are 5 unpaired electrons and for Fe^{2+} , there are 4 unpaired electrons, leading to magnetic moments of 5.92 and 4.90 B.M., respectively.

Quick Tip

For calculating the spin-only magnetic moment, determine the number of unpaired electrons using the electronic configuration of the metal in the complex.

52. For hydrogen-like species, which of the following graphs provides the most appropriate representation of E vs Z plot for a constant n ?



Correct Answer: (2)

Solution: For hydrogen-like species, the energy of the stationary state is proportional to Z^2 , where Z is the atomic number. Hence, the correct plot for E vs Z is a quadratic curve, showing $E \propto Z^2$.

Quick Tip

The energy of a hydrogen-like atom is proportional to Z^2 , so the plot of E versus Z should show a parabolic increase.

53. Given below are two statements:

Statement (I): In partition chromatography, the stationary phase is a thin film of liquid present in the inert support.

Statement (II): In paper chromatography, the material of paper acts as a stationary phase.

In light of the above statements, choose the correct answer from the options given below:

- (1) Both Statement I and Statement II are false
- (2) Statement I is true but Statement II is false
- (3) Both Statement I and Statement II are true
- (4) Statement I is false but Statement II is true

Correct Answer: (2)

Solution: In partition chromatography, the stationary phase is indeed a thin liquid film that is supported by an inert material, so Statement I is true. However, in paper chromatography, the stationary phase is actually water present in the paper fibers, not the material of the paper itself, so Statement II is false.

Quick Tip

In partition chromatography, the stationary phase is a liquid film, while in paper chromatography, water acts as the stationary phase.

54. Identify the essential amino acids from below:

(A) Valine (B) Proline (C) Lysine (D) Threonine (E) Tyrosine

Choose the correct answer from the options given below:

- (1) (A), (C) and (D) only
- (2) (A), (C) and (E) only

(3) (B), (C) and (E) only

(4) (C), (D) and (E) only

Correct Answer: (1) (A), (C) and (D) only

Solution:

Essential amino acids are those that cannot be synthesized by the human body and must be obtained through the diet. Among the amino acids listed:

- **Valine (A)** is an essential amino acid.

- **Proline (B)** is a non-essential amino acid, as the body can synthesize it.

- **Lysine (C)** is an essential amino acid.

- **Threonine (D)** is an essential amino acid.



- **Tyrosine (E)** is a non-essential amino acid because it can be synthesized from phenylalanine, which is essential.

Thus, the essential amino acids are Valine, Lysine, and Threonine, making the correct answer (1).

Quick Tip

Essential amino acids are those that the body cannot produce on its own, and must be acquired from food. Examples include Valine, Lysine, and Threonine.

55. Which among the following halides will generate the most stable carbocation in a nucleophilic substitution reaction?



Correct Answer: (4)

Solution:

The stability of carbocations depends on the extent of delocalization and resonance stabilization. In the case of the halides:


- The carbocation generated from option (4) ($\text{C}_6\text{H}_5\text{C}(\text{Br}) - \text{Ph}$) is the most stable because it benefits from resonance stabilization provided by the aromatic phenyl group.
- The carbocations formed in options (1), (2), and (3) are less stable, as they lack this additional resonance stabilization.

Thus, the most stable carbocation is generated from option (4).

Quick Tip

Resonance stabilization from aromatic rings significantly increases the stability of carbocations. Therefore, halides that lead to resonance-stabilized carbocations tend to undergo nucleophilic substitution reactions more readily.

56. Consider the equilibrium:



If the pressure applied over the system increases by two fold at constant temperature then:

- (1) (A) and (B) only
- (2) (A), (B) and (D) only
- (3) (B) and (C) only
- (4) (A), (B) and (C) only

Correct Answer: (1) (A) and (B) only

Solution:

According to Le Chatelier's principle, increasing the pressure on a gaseous system will favor the side with fewer moles of gas. In the given equilibrium reaction, the left-hand side has 4 moles of gas (1 mole of CO and 3 moles of H_2) and the right-hand side has 2 moles of gas (1 mole of CH_4 and 1 mole of H_2O). Therefore, increasing the pressure will shift the equilibrium towards the right (in the forward direction), increasing the concentration of products and decreasing the concentration of reactants.

- **(A)** The concentration of reactants and products increases because the equilibrium shifts toward the products side.
- **(B)** The equilibrium will shift in the forward direction to produce more CH_4 and H_2O .

- (C) The equilibrium constant remains unchanged, as pressure does not affect the value of the equilibrium constant at constant temperature.

Therefore, the correct answer is (1) (A) and (B) only.

Quick Tip

Le Chatelier's principle helps predict the direction of equilibrium shifts when a system is disturbed. Increasing pressure favors the side with fewer gas molecules in a reaction.

57. Given below are two statements:

Statement (I): NaCl is added to the ice at 0°C , present in the ice cream box to prevent the melting of ice cream.

Statement (II): On addition of NaCl to ice at 0°C , there is a depression in freezing point.

In the light of the above statements, choose the correct answer from the options given below:

- (1) Statement I is false but Statement II is true
- (2) Both Statement I and Statement II are true
- (3) Statement I is false but Statement II is true
- (4) Statement I is true but Statement II is false

Correct Answer: (2) Both Statement I and Statement II are true

Solution:

- **Statement (I):** The addition of NaCl to ice helps lower the freezing point of water, preventing the ice from melting at 0°C and thus allowing for the ice cream to stay frozen at lower temperatures. This statement is correct.

- **Statement (II):** Adding NaCl to ice creates a phenomenon known as freezing point depression, which lowers the freezing point of water and ice. This is a well-known colligative property of solutions. Hence, this statement is also correct.

Therefore, both statements are true.

Quick Tip

Freezing point depression occurs when a solute (like NaCl) is added to a solvent (like water), causing the freezing point to lower. This is a colligative property that depends on the concentration of solute particles.

58. Given below are two statements:

Statement (I): On nitration of m-xylene with HNO_3 , H_2SO_4 , followed by oxidation, 4-nitrobenzene-1, 3-dicarboxylic acid is obtained as the major product.

Statement (II): CH_3 group is o/p-directing while NO_2 group is m-directing group.

In the light of the above statements, choose the correct answer from the options given below:

- (1) Statement I is false but Statement II is true
- (2) Statement I is false but Statement II is true
- (3) Both Statement I and Statement II are true
- (4) Statement I is true but Statement II is false

Correct Answer: (3) Both Statement I and Statement II are true

Solution:

- **Statement (I):** The nitration of m-xylene with HNO_3 and H_2SO_4 followed by oxidation indeed gives 4-nitrobenzene-1,3-dicarboxylic acid as the major product. This statement is correct.

- **Statement (II):** The methyl group (CH_3) is o/p-directing, meaning it directs electrophilic substitution reactions to the ortho and para positions. The nitro group (NO_2) is m-directing, meaning it directs electrophilic substitution to the meta position. This is also correct.

Therefore, both statements are true.

Quick Tip

The methyl group (CH_3) is an electron-donating group that activates the aromatic ring towards electrophilic substitution at the o/p-positions. The nitro group (NO_2) is an electron-withdrawing group that directs substitution to the meta-position.

59. 0.1 M solution of KI reacts with excess of H₂SO₄ and KIO₃, according to the equation:



Identify the correct statements: (A) 200 mL of KI solution reacts with 0.004 mol of KIO₃ (B) 200 mL of KI solution reacts with 0.006 mol of H₂SO₄ (C) 0.5 L of KI solution produced 0.005 mol of I₂ (D) Equivalent weight of KIO₃ is equal to:

$$\frac{\text{Molecular weight}}{5}$$

Choose the correct answer from the options given below:

- (1) (A) and (D) only
- (2) (A) and (B) only
- (3) (B) and (C) only
- (4) (C) and (D) only

Correct Answer: (1) (A) and (D) only

Solution:

- Statement (A) is correct because 200 mL of 0.1 M KI contains 0.02 moles of KI, and according to the equation, 5 moles of I⁻ react with 1 mole of KIO₃. Therefore, 0.02 moles of KI would require 0.004 mol of KIO₃.
- Statement (C) is also correct because 0.5 L of 0.1 M KI will contain 0.05 moles of KI, and according to the equation, this will produce 0.005 mol of I₂.
- Statement (D) is correct because the equivalent weight of KIO₃ is equal to its molecular weight divided by 5, as 5 moles of iodide react with one mole of KIO₃.

Thus, the correct answer is (A), (D).

Quick Tip

The equivalent weight of a substance is defined as its molecular weight divided by the number of electrons involved in the reaction. For KIO₃, 5 moles of electrons are involved per mole of KIO₃.

60. Match List-I with List-II:

List-I Applications		List-II Batteries/Cell	
(A)	Transistors	(I)	Anode - Zn/Hg ; Cathode - HgO + C
(B)	Hearing aids	(II)	Hydrogen fuel cell
(C)	Invertors	(III)	Anode - Zn; Cathode - Carbon
(D)	Apollo space ship	(IV)	Anode - Pb ; Cathode - Pb PbO ₂

Choose the correct answer from the options given below:

- (1) (A)-(III), (B)-(I), (C)-(IV), (D)-(II)
- (2) (A)-(III), (B)-(II), (C)-(I), (D)-(IV)
- (3) (A)-(II), (B)-(III), (C)-(IV), (D)-(I)
- (4) (A)-(III), (B)-(III), (C)-(IV), (D)-(I)

Correct Answer: (1) (A)-(III), (B)-(I), (C)-(IV), (D)-(II)

Solution:

- **Transistors (A)** use a battery with an anode of Zn/Hg and a cathode of HgO + C.
- **Hearing aids (B)** use a battery with an anode of Zn and a cathode of Carbon.
- **Invertors (C)** use a hydrogen fuel cell as the energy source.
- **Apollo space ship (D)** uses a battery with an anode of Pb and a cathode of PbO₂.

Therefore, the correct matching is $(A) - (III), (B) - (I), (C) - (IV), (D) - (II)$.

Quick Tip

Different applications require different types of batteries depending on the energy needs and the chemical reactions involved in generating electricity.

61. O₂ gas will be evolved as a product of electrolysis of:

- (A) an aqueous solution of AgNO₃ using silver electrodes.
- (B) an aqueous solution of AgNO₃ using platinum electrodes.
- (C) a dilute solution of H₂SO₄ using platinum electrodes.
- (D) a high concentration solution of H₂SO₄ using platinum electrodes.

Choose the correct answer from the options given below:

- (1) (B) and (C) only
- (2) (A) and (C) only
- (3) (B) and (D) only
- (4) (A) and (C) only

Correct Answer: (1) (B) and (C) only

Solution:

In the electrolysis of water or aqueous solutions, oxygen gas (O_2) is produced at the anode under certain conditions.

- **(A) An aqueous solution of $AgNO_3$ using silver electrodes:** In this case, silver ions are reduced at the cathode to form silver metal. Oxygen gas is not produced at the anode because the silver electrode undergoes oxidation to form silver ions. Therefore, no O_2 is evolved.

- **(B) An aqueous solution of $AgNO_3$ using platinum electrodes:** Platinum is an inert electrode, and when $AgNO_3$ is electrolyzed, oxygen gas will evolve at the anode due to the oxidation of water. This is correct.

- **(C) A dilute solution of H_2SO_4 using platinum electrodes:** In this case, water is the main electrolyte, and oxygen gas will evolve at the anode during electrolysis of the dilute sulfuric acid solution. This is correct.

- **(D) A high concentration solution of H_2SO_4 using platinum electrodes:** At high concentrations of H_2SO_4 , oxygen evolution is suppressed, and hydrogen gas is more likely to evolve at the anode. This is not correct for oxygen evolution.


Thus, the correct answer is **(1)** (B) and (C) only.

Quick Tip

Oxygen gas is evolved at the anode during electrolysis when water or dilute sulfuric acid is electrolyzed. At high concentrations of sulfuric acid, hydrogen gas evolution is favored at the anode.

62. Identify the homoleptic complexes with odd number of d-electrons in the central metal:

- (A) $[FeO_4]^{2-}$
- (B) $[Fe(CN)_6]^{3-}$



Choose the correct answer from the options given below:

(1) (B) and (D) only

(2) (A), (B) and (D) only

(3) (A), (B) and (E) only

(4) (A), (C), (D) and (E) only

Correct Answer: (1) (B) and (D) only

Solution:

- (A) $[FeO_4]^{2-}$: Iron in this complex is in the +2 oxidation state, with d^6 electrons. Since 6 is an even number, this complex does not meet the requirement for an odd number of d-electrons.

- (B) $[Fe(CN)_6]^{3-}$: Iron in this complex is in the +3 oxidation state, with d^5 electrons. This gives an odd number of d-electrons, so this is a homoleptic complex with odd d-electrons.

- (C) $[Fe(CN)_6]^{2-}$: Iron is in the +2 oxidation state with d^6 electrons. This complex has an even number of d-electrons.

- (D) $[CoCl_4]^{2-}$: Cobalt in this complex is in the +2 oxidation state, with d^7 electrons. This gives an odd number of d-electrons, so this is a homoleptic complex with odd d-electrons.

- (E) $[Co(H_2O)_6]^{3+}$: Cobalt in the +3 oxidation state has d^6 electrons, which is an even number.

Thus, the correct answer is (1) (B) and (D) only.

Quick Tip

To identify complexes with odd numbers of d-electrons, determine the oxidation state of the central metal and subtract the oxidation state from the total number of electrons in the neutral atom.

63. Total number of sigma (σ) and pi (π) bonds respectively present in hex-1-en-4-yne are:

(1) 13 and 3

- (2) 11 and 3
(3) 13 and 13
(4) 14 and 3

Correct Answer: (1) 13 and 3

Solution:

The structure of hex-1-en-4-yne consists of:

- A C=C double bond (with one σ -bond and one π -bond).
- A CC triple bond (with one σ -bond and two π -bonds).
- Six single bonds between carbon atoms and hydrogen atoms (each σ -bonds).

Count of σ -bonds:

- 1 σ -bond in the C=C double bond.
- 1 σ -bond in the CC triple bond.
- 11 σ -bonds between carbon and hydrogen atoms.

Thus, the total number of σ -bonds is 13.

Count of π -bonds:

- 1 π -bond in the C=C double bond.
- 2 π -bonds in the CC triple bond.



Thus, the total number of π -bonds is 3.

Therefore, the correct answer is **(1)** 13 and 3.

Quick Tip

When counting bonds in organic compounds, remember that each single bond is a σ -bond, each double bond consists of one σ -bond and one π -bond, and each triple bond consists of one σ -bond and two π -bonds.

64. If



At constant temperature. Then:

(1) $X = Y + Z$

(2) $X - Y = Z$

(3) $X = Y - Z$

(4) $X = Y + Z$

Correct Answer: (4) $X = Y + Z$

Solution:

The energy change when diamond converts to graphite is X . The total enthalpy change for the complete oxidation of diamond to carbon dioxide is the sum of the enthalpy changes of the oxidation steps of both diamond and graphite. Therefore, the enthalpy change for the overall reaction is the sum of Y (oxidation of diamond) and Z (oxidation of graphite).

Hence, $X = Y + Z$.

Thus, the correct answer is (4) $X = Y + Z$.

Quick Tip

The enthalpy changes for reactions can be combined if the reactions occur sequentially, such as in the oxidation of diamond and graphite to form carbon dioxide.

65. Given below are two statements:

Statement (I): It is impossible to specify simultaneously with arbitrary precision, the linear momentum and the position of a particle.

Statement (II): If the uncertainty in the measurement of position and uncertainty in measurement of momentum are equal for an electron, then the uncertainty in the measurement of velocity is $\geq \sqrt{\frac{h}{\pi}} \times \frac{1}{2m}$.

In the light of the above statements, choose the correct answer from the options given below:

(1) Statement I is true but Statement II is false.

(2) Both Statement I and Statement II are true.

(3) Statement I is false but Statement II is true.

(4) Both Statement I and Statement II are false.

Correct Answer: (2) Both Statement I and Statement II are true.

Solution:

- **Statement I:** This is a restatement of the Heisenberg Uncertainty Principle, which asserts that it is impossible to precisely determine both the position and the momentum of a particle simultaneously. This statement is true.

- **Statement II:** The Heisenberg Uncertainty Principle provides the relationship between the uncertainty in position (Δx) and momentum (Δp), given by:

$$\Delta x \Delta p \geq \frac{h}{4\pi}$$

For an electron, if the uncertainties in position and momentum are equal, the uncertainty in velocity Δv can be expressed as:

$$\Delta v = \frac{\Delta p}{m} \geq \sqrt{\frac{h}{\pi}} \times \frac{1}{2m}$$






This statement is also correct.

Therefore, the correct answer is (2) Both Statement I and Statement II are true.

Quick Tip

The Heisenberg Uncertainty Principle is fundamental in quantum mechanics, stating that we cannot simultaneously know the exact position and momentum of a particle. The uncertainty in velocity is derived from the uncertainties in position and momentum.


66. Which one of the following reaction sequences will give an azo dye?



Solution:

An azo dye is formed through a reaction called the diazotization reaction, where a primary aromatic amine reacts with sodium nitrite in an acidic medium, forming a diazonium salt.

This salt then reacts with an aromatic compound, such as naphthol, to form an azo dye.



The correct reaction sequence for the formation of an azo dye involves the following steps:

1. NO₂ (i) Sn/HCl reduces the nitro group (NO₂) to an amine group (NH₂).
2. (ii) NaNO₂/HCl diazotizes the amine group to form a diazonium salt (Ar-N₂⁺).
3. (iii) β-naphthol, NaOH reacts with the diazonium salt to form the azo dye.

Thus, the correct reaction sequence that will give an azo dye is option **(1)**.

Quick Tip

The formation of azo dyes involves the diazotization of aromatic amines followed by coupling with compounds like naphthols. This is an important reaction in dye chemistry.

67. Drug X becomes ineffective after 50% decomposition. The original concentration of drug in a bottle was 16 mg/mL which becomes 4 mg/mL in 12 months. The expiry time of the drug in months is ____ . Assume that the decomposition of the drug follows first order kinetics.

- (1) 12
- (2) 2
- (3) 3
- (4) 6

Correct Answer: (4) 6

Solution:

For a first-order reaction, the equation for the concentration at time t is given by:

$$\ln \left(\frac{[A]_0}{[A]_t} \right) = kt$$

where $[A]_0$ is the initial concentration, $[A]_t$ is the concentration at time t , k is the rate constant, and t is the time.

Given that the drug becomes ineffective after 50% decomposition, we know that

$[A]_0 = 16 \text{ mg/mL}$ and $[A]_t = 4 \text{ mg/mL}$. Substitute the values into the equation:

$$\ln \left(\frac{16}{4} \right) = k(12)$$

$$\ln(4) = 12k \quad \Rightarrow \quad 1.386 = 12k \quad \Rightarrow \quad k = 0.1155 \text{ month}^{-1}$$

Now, to find the expiry time, we use the same equation for 50% decomposition (i.e.,

$[A]_t = \frac{[A]_0}{2}$):

$$\ln \left(\frac{[A]_0}{\frac{[A]_0}{2}} \right) = k \cdot t_{\text{expiry}}$$

$$\ln(2) = 0.1155 \cdot t_{\text{expiry}} \quad \Rightarrow \quad 0.693 = 0.1155 \cdot t_{\text{expiry}} \quad \Rightarrow \quad t_{\text{expiry}} = 6 \text{ months}$$

Thus, the expiry time of the drug is 6 months.

Quick Tip

For first-order reactions, the time for 50% decomposition can be calculated using the equation: $\ln \left(\frac{[A]_0}{[A]_t} \right) = kt$. The rate constant can be used to calculate the time for any given concentration change.

68. The type of oxide formed by the element among Li, Na, Be, Mg, B and Al that has the least atomic radius is:

- (1) A_2O_3
- (2) AO_2
- (3) A_2O
- (4) A_2O_4

Correct Answer: (1) A_2O_3

Solution:

The atomic radius generally decreases across a period from left to right due to increasing nuclear charge. Among the elements given (Li, Na, Be, Mg, B, Al), the atomic radius decreases as we move across the period from Li to Al. The element that forms the oxide with the least atomic radius is Al (aluminum), which forms Al_2O_3 .

Thus, the correct answer is .

Quick Tip

The atomic radius decreases across a period from left to right due to an increase in nuclear charge, which pulls the electrons closer to the nucleus.

69. First ionisation enthalpy values of the first four group 15 elements are given below. Choose the correct value for the element that is a main component of the apatite family:

- (1) 1012 kJ mol^{-1}
- (2) 1402 kJ mol^{-1}
- (3) 834 kJ mol^{-1}
- (4) 947 kJ mol^{-1}

Correct Answer: (1) 1012 kJ mol^{-1}

Solution:


The apatite family is composed of minerals that are mainly made of phosphate ions, and one of the key elements in this family is phosphorus. Phosphorus is the element in Group 15, and its ionization enthalpy is 1012 kJ mol^{-1} . Hence, the correct value corresponds to the ionization enthalpy of phosphorus.

Thus, the correct answer is .

Quick Tip

The ionization enthalpy increases across a period as the effective nuclear charge increases, but it decreases down a group due to an increase in atomic size. Phosphorus is a key element in the apatite family and has a relatively moderate ionization enthalpy.

70. Which one of the following, with HBr, will give a phenol?



Correct Answer: (2) $\text{CH}_3\text{C}_6\text{H}_4\text{OCH}_3$

Solution:

The reaction of an aromatic ether with HBr results in the cleavage of the C-O bond, leading to the formation of a phenol. For the compound $\text{CH}_3\text{C}_6\text{H}_4\text{OCH}_3$, this cleavage results in the formation of phenol ($\text{C}_6\text{H}_5\text{OH}$) and methyl bromide (CH_3Br).

Thus, the correct answer is .

Quick Tip

Aromatic ethers react with HBr to undergo nucleophilic substitution, where the bond between the oxygen atom and the alkyl group is broken, yielding phenol and the corresponding alkyl halide.

(Chemistry) Section-B

71. Consider the following low-spin complexes



The sum of the spin-only magnetic moment values of complexes having yellow colour is:

B.M. (answer is nearest integer)

Solution:

To determine the magnetic moment, we first need to consider the electronic configuration of the metal ions and their respective spin states. We are looking for the complexes with yellow colour, which are low-spin complexes.

- (I) $\text{K}_3[\text{Co}(\text{NO}_3)_6]$:

- In the presence of a ligand field, Co^{3+} (d^6 configuration) undergoes d^2sp^3 hybridization, and it is a low-spin complex. - Magnetic moment:

$$\mu = \sqrt{n(n+2)} = 0 \text{ B.M.}$$

- (II) $\text{K}_4[\text{Fe}(\text{CN})_6]$:

- Fe^{2+} (d^6 configuration) undergoes d^2sp^3 hybridization, and it is also a low-spin complex. - Magnetic moment:

$$\mu = \sqrt{n(n+2)} = 0 \text{ B.M.}$$

Thus, the sum of the spin-only magnetic moment values for the complexes with yellow colour is 0 B.M..

Therefore, the correct answer is $\boxed{(0)}$.

Quick Tip

For low-spin complexes, the metal ion undergoes hybridization that results in paired electrons in the lower energy orbitals, leading to no unpaired electrons and hence a magnetic moment of 0 B.M.

72. Isomeric hydrocarbons \rightarrow negative Baeyer's test (Molecular formula C_9H_{12}). The total number of isomers from above with four different non-aliphatic substitution sites is -

Solution:

The molecular formula C_9H_{12} indicates that the compound is likely to have several isomers due to the possibility of different types of substitutions. For this molecular formula, there can be isomers with various positions for non-aliphatic substitution.

There are 3 distinct isomers of C_9H_{12} with four different non-aliphatic substitution sites as shown in the answer. These include positional isomers and functional group isomers.

Thus, the correct number of isomers is $\boxed{(2)}$.

Quick Tip

When dealing with isomeric hydrocarbons, focus on the positions of non-aliphatic groups and the potential for different bonding arrangements.

73. In the Claisen-Schmidt reaction to prepare dibenzalacetone from 5.3 g benzaldehyde, a total of 3.51 g of product was obtained. The percentage yield in this reaction was _____.

Solution:

First, calculate the moles of benzaldehyde used:

$$\text{Moles of benzaldehyde} = \frac{5.3}{106} = 0.05 \text{ mol}$$

The theoretical yield of the product can be calculated from the stoichiometry of the reaction (based on a 1:1 molar ratio for reactants and products):

$$\text{Theoretical yield} = 0.05 \text{ mol} \times \frac{233 \text{ g/mol}}{1} = 11.65 \text{ g}$$

Now, calculate the percentage yield:

$$\% \text{yield} = \frac{3.51 \text{ g}}{11.65 \text{ g}} \times 100 = 30\%$$

Thus, the percentage yield of the reaction is 60.

Quick Tip

The percentage yield can be calculated by comparing the actual yield to the theoretical yield, using the formula:

$$\% \text{yield} = \frac{\text{actual yield}}{\text{theoretical yield}} \times 100$$

74. In the sulphur estimation, 0.20 g of a pure organic compound gave 0.40 g of barium sulphate. The percentage of sulphur in the compound is

Solution:

Given that the molar mass of sulphur (S) is 32 g/mol and that of barium sulphate (BaSO₄) is 233 g/mol, the mass of sulphur in the compound can be calculated from the mass of barium sulphate produced:

The moles of barium sulphate formed:

$$\text{Moles of BaSO}_4 = \frac{0.40 \text{ g}}{233 \text{ g/mol}} = 0.00172 \text{ mol}$$

The moles of sulphur in the compound are equal to the moles of BaSO₄ because of the 1:1 stoichiometry of BaSO₄ and sulphur.

The mass of sulphur is:

$$\text{Mass of S} = 0.00172 \text{ mol} \times 32 \text{ g/mol} = 0.05504 \text{ g}$$

The percentage of sulphur in the compound is:

$$\%S = \frac{0.05504 \text{ g}}{0.20 \text{ g}} \times 100 = 27.5\%$$

Thus, the correct percentage of sulphur is 27.5%.

Quick Tip

To find the percentage of an element in a compound, divide the mass of the element by the mass of the sample and multiply by 100.

75. Total number of non-bonded electrons present in NO₂⁻ ion based on Lewis theory is:

- (1) 10
- (2) 12
- (3) 8
- (4) 6

Correct Answer: (12)

Solution:

For the NO₂⁻ ion, we consider the electron configuration based on the Lewis structure. The total number of electrons in NO₂⁻ is the sum of the electrons from nitrogen and oxygen atoms. Nitrogen has 5 valence electrons and each oxygen has 6 valence electrons. The negative charge adds an additional electron.

The total number of valence electrons is:

$$5 \text{ (from N)} + 6 \times 2 \text{ (from 2 O)} + 1 \text{ (from the negative charge)} = 18 \text{ electrons}$$

In the Lewis structure of NO_2^- , 2 electrons are involved in the bonding between nitrogen and oxygen atoms, while the remaining electrons are non-bonded.

Thus, the number of non-bonded electrons is .

Quick Tip

In Lewis structures, non-bonded electrons are the valence electrons that are not shared between atoms. Count these electrons after assigning bonds to the atoms.