

## Syllabus for TG EAPCET 2025-E Stream (Engineering Stream)

### MODEL QUESTIONS – MATHEMATICS

- 1) I) The coefficient of  $x^5$  in  $(1 - x - x^2 + x^3)^6$  is 20.  
II) If p and q are the coefficients of  $x^4$  in  $(1 + x)^{2n}$  and  $(1 + x)^{2n-1}$  respectively then  $2p=q$ .  
Which of the above statements is (are) true.
- 1) only I                      2) Only II                      3) Both I and II                      4) Neither I nor II
- 2). Assertion (A): If  $A+B=45^\circ$  then  $(1+\tan A)(1+\tan B)=2$ .  
Reason(R):  $\tan 22\frac{1}{2}^\circ = \sqrt{2} - 1$ .
- 1) Both A & R are True and R is the correct explanation of A.  
2) Both A & R are True and R is not correct explanation of A.  
3) A is True but R is False.  
4) R is True but A is False.
- 3) Arrange the following statements in ascending order of their results
- A) The order of  $\left(\frac{d^4y}{dx^4} + \frac{d^2y}{dx^2}\right)^{3/2} = a\left(\frac{d^3y}{dx^3}\right)$   
B) The degree of  $\left(\frac{dy}{dx} + \frac{d^2y}{dx^2}\right)^{5/4} = a\left(\frac{d^3y}{dx^3}\right)^{2/3}$   
C) The degree of  $y = \left(1 + \left(\frac{dy}{dx}\right)^2\right)^{3/2} \left(\frac{d^2y}{dx^2}\right)$   
D) The order of  $x^3 + \left(\frac{d^3y}{dx^3}\right)^2 + 2x^2\left(\frac{d^2y}{dx^2}\right) - 3y = x^2$
- 1) D,C,B,A                      2) D,C,A,B                      3) C,D,B,A                      4) C,D,A,B
- 4) A and B are two independent events of a sample space such that  $P(A)=0.2$ ,  $P(B)=0.5$ .
- | List I           | List II  |
|------------------|----------|
| A) $P(B/A)$      | I) 0.2   |
| B) $P(A/B)$      | II) 0.1  |
| C) $P(A \cap B)$ | III) 0.3 |
| D) $P(A \cup B)$ | IV) 0.6  |
|                  | V) 0.5   |
- The correct match is
- 1) A-IV, B-V, C-III, D-I  
2) A-V, B-I, C-II, D-IV  
3) A-III, B-I, C-II, D-IV  
4) A-II, B-I, C-IV, D-V
- 5) The line  $ax + by + c = 0$  is a normal to the circle  $x^2 + y^2 + 4x + 6y + 8 = 0$  if and only if
- (1)  $2a + 3b = c$     (2)  $3a + 2b = c$     (3)  $2a + 3b + c = 0$     (4)  $3a + 2b + c = 0$

## Syllabus for TG EAPCET 2025-E Stream (Engineering Stream)

- 6)  $\begin{vmatrix} al & bm & cn \\ l^2 & m^2 & n^2 \\ 1 & 1 & 1 \end{vmatrix} = \Delta_1$  and  $\begin{vmatrix} a & b & c \\ l & m & n \\ mn & ln & lm \end{vmatrix} = \Delta_2$  then
- (1)  $\Delta_1 = \Delta_2$                       (2)  $\Delta_1 = 2\Delta_2$                       (3)  $2\Delta_1 = \Delta_2$                       (4)  $\Delta_1 + \Delta_2 = 0$
- 7) If  $\vec{a}$  is a non-zero vector and  $\vec{b}, \vec{c}$  are two vectors such that  $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$  and  $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$  then
- 1)  $\vec{b} - \vec{c}$  is collinear with  $\vec{a}$                       2)  $\vec{b} - \vec{c}$  is perpendicular with  $\vec{a}$   
 3)  $\vec{b} = \vec{c}$     4)  $\vec{b} \neq \vec{c}$
- 8) If  $L_1 : 2x + 3y - 20 = 0$ ,  $L_2 : 2x + 3y - 14 = 0$ , then the straight line represented by  $a(2x + 3y - 20) + b(2x + 3y - 14) = 0$  is
- 1) Parallel to  $L_1 = 0$  and  $L_2 = 0$ .  
 2) Perpendicular to  $L_1 = 0$  and parallel to  $L_2 = 0$   
 3) Perpendicular to  $L_1 = 0$  and  $L_2 = 0$   
 4) Parallel to  $L_1 = 0$  and Perpendicular to  $L_2 = 0$
- 9) If  $f(x) = \log\left(\frac{1+2x}{1-2x}\right)$ , then  $x =$
- 1)  $\frac{e^{f(x)} - 1}{2(e^{f(x)} + 1)}$                       2)  $\frac{2(e^{f(x)} - 1)}{(e^{f(x)} + 1)}$                       3)  $\frac{e^{f(x)} - 1}{(e^{f(x)} + 1)}$                       4)  $\frac{e^{f(2x)} - 1}{(e^{f(2x)} + 1)}$
- 10) If  $\tan\theta = \frac{b}{a}$  then  $a \cos 2\theta - b \sin 2\theta =$
- 1)  $a$     2)  $b$     3)  $2a$     4)  $2b$

\*\*\*\*\*