

# Limit, Continuity, And Differentiability JEE Main PYQ – 3

Total Time: 25 Minute

Total Marks: 40

## Instructions

### Instructions

1. Test will auto submit when the Time is up.
2. The Test comprises of multiple choice questions (MCQ) with one or more correct answers.
3. The clock in the top right corner will display the remaining time available for you to complete the examination.

### Navigating & Answering a Question

1. The answer will be saved automatically upon clicking on an option amongst the given choices of answer.
2. To deselect your chosen answer, click on the clear response button.
3. The marking scheme will be displayed for each question on the top right corner of the test window.

## Limit, Continuity, And Differentiability

1. If  $f(x) = [a+13 \sin x]$  &  $x \in (0, \pi)$ , then number of non-differentiable points of  $f(x)$  are [where 'a' is integer] **(+4, -1)**
- 
2. If  $f$  and  $g$  are differentiable functions in  $(0, 1)$  satisfying  $f(0) = 2 = g(1), g(0) = 0$  and  $f(1) = 6$ , then for some  $c \in ]0, 1[$  **(+4, -1)**
- a.  $2f'(c) = g'(c)$
- b.  $2f'(c) = 3g'(c)$
- c.  $f'(c) = g'(c)$
- d.  $f'(c) = 2g'(c)$
- 
3. For  $x \in \mathbb{R}, f(x) = |\log 2 - \sin x|$  and  $g(x) = f(f(x))$ , then : **(+4, -1)**
- a.  $g$  is not differentiable at  $x = 0$
- b.  $g'(0) = \cos(\log 2)$
- c.  $g'(0) = -\cos(\log 2)$
- d.  $g$  is differentiable at  $x = 0$  and  $g'(0) = -\sin(\log 2)$
- 
4. If for  $x \in (0, \frac{1}{4})$ , the derivative of  $\tan^{-1} \left( \frac{6x\sqrt{x}}{1-9x^3} \right)$  is  $\sqrt{x} \cdot g(x)$ , then  $g(x)$  equals : **(+4, -1)**
- a.  $\frac{3x\sqrt{x}}{1-9x^3}$
- b.  $\frac{3x}{1-9x^3}$
- c.  $\frac{3}{1+9x^3}$
- d.  $\frac{9}{1+9x^3}$
- 
5. For  $x > 1$ , if  $(2x)^{2y} = 4e^{2x-2y}$ , then  $(1 + \log_e 2x)^2 \frac{dy}{dx}$  is equal to : **(+4, -1)**

- a.  $\log_e 2x$
- b.  $\frac{x \log_e 2x + \log_e 2}{x}$
- c.  $x \log_e 2x$
- d.  $\frac{x \log_e 2x - \log_e 2}{x}$

---

6. If  $f(x) = \sin(\sin x)$  and  $f''(x) + \tan x f'(x) + g(x) = 0$ , then  $g(x)$  is : (+4, -1)

- a.  $\cos^2 x \cos(\sin x)$
- b.  $\sin^2 x \cos(\cos x)$
- c.  $\sin^2 x \sin(\cos x)$
- d.  $\cos^2 x \sin(\sin x)$

---

7. If  $x = \sqrt{2^{\operatorname{cosec}^{-1} t}}$  and  $y = \sqrt{2^{\operatorname{sec}^{-1} t}} (|t| \geq 1)$ , then  $\frac{dy}{dx}$  is equal to : (+4, -1)

- a.  $\frac{y}{x}$
- b.  $\frac{x}{y}$
- c.  $-\frac{y}{x}$
- d.  $-\frac{x}{y}$

---

8. If  $x = 3 \tan t$  and  $y = 3 \sec t$ , then the value of  $\frac{d^2y}{dx^2}$  at  $t = \frac{\pi}{4}$ , is : (+4, -1)

- a.  $\frac{3}{2\sqrt{2}}$
  - b.  $\frac{1}{3\sqrt{2}}$
  - c.  $\frac{1}{6}$
  - d.  $\frac{1}{6\sqrt{2}}$
-

9. Let  $f : (-1, 1) \rightarrow \mathbb{R}$  be a function defined by  $f(x) = \max\{-|x|, -\sqrt{1-x^2}\}$ . If  $K$  be the set of all points at which  $f$  is not differentiable, then  $K$  has exactly : **(+4, -1)**

- a. Three elements
- b. One element
- c. Five elements
- d. Two elements

---

10. Let  $k$  be a non-zero real number. If  $f(x) = \begin{cases} \frac{\left(e^{x-1}\right)^2 \sin\left(\frac{x}{k}\right) \log\left(1+\frac{x}{4}\right), & \text{if } x \neq 0 \\ \sqrt[2]{e^x} \log 2, & \text{if } x = 0 \end{cases}$  is a continuous function, then the value of  $k$  is : **(+4, -1)**

- a. 1
- b. 2
- c. 3
- d. 4




## Answers

1. Answer: 25 – 25

Explanation:

Points where  $\sin X = \frac{1}{3}, \frac{2}{13}, \dots, \frac{12}{13}$  will be the points of non-derivability of  $f(x)$   
 $\Rightarrow$  24 points



and also where the  $\sin x = 1 \Rightarrow$  1 points.  
 Therefore, 25 points of non-derivability

Concepts:

1. Continuity & Differentiability:

### Definition of Differentiability

$f(x)$  is said to be differentiable at the point  $x = a$ , if the derivative  $f'(a)$  be at every point in its domain. It is given by

The maxima lie between the minima and the width of the central maximum is simply the distance between the 1st order minima from the centre of the screen on both sides of the centre.

The position of the minima given by  $y$  (measured from the centre of the screen) is:

$$\tan\theta \approx \theta = y/D$$

For small  $\theta$ ,

$$\sin \theta \approx \theta$$

$$\Rightarrow \lambda = a \sin \theta \approx a\theta$$

$$\Rightarrow \theta = y/D = \lambda/a$$

$$\Rightarrow y = \lambda Da$$

The width of the central maximum is simply twice this value

$$\Rightarrow \text{Width of central maximum} = 2\lambda Da$$

$$\Rightarrow \text{Angular width of central maximum} = 2\theta = 2\lambda/a$$

## Definition of Continuity

Mathematically, a function is said to be continuous at a point  $x = a$ , if

It is implicit that if the left-hand limit (L.H.L), right-hand limit (R.H.L), and the value of the function at  $x=a$  exist and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

$$\lim_{x \rightarrow a} f(x) \text{ Exists, and}$$

$$\lim_{x \rightarrow a} f(x) = f(a)$$

If the function is unspecified or does not exist, then we say that the function is discontinuous.

### 2. Answer: d

#### Explanation:

Using, mean value theorem

$$f'(c) = \frac{f(1)-f(0)}{1-0} = 4$$

$$g'(c) = \frac{g(1)-g(0)}{1-0} = 2$$

$$\text{so, } f'(c) = 2g'(c)$$

#### Concepts:

## 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

---

## 3. Answer: b

### Explanation:

$$g(x) = |\log_e 2 - \sin(|\log_e 2 - \sin x|)|$$

$$\text{At } x = 0, g(x) = \log_e(2) - \sin(\log_e 2 - \sin x)$$

$$\therefore g'(x) = \cos(\log_e(2) - \sin x) \times \cos(x)$$

$$\Rightarrow g'(0) = \cos(\log_e(2))$$

### Concepts:

## 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

---

#### 4. Answer: d

**Explanation:**

$$f(x) = 2 \tan^{-1}(3x\sqrt{x})$$

$$\text{For } x \in \left(0, \frac{1}{4}\right)$$

$$f'(x) = \frac{9\sqrt{x}}{1+9x^3}$$

$$g(x) = \frac{9}{1+9x^3}$$

**Concepts:**

##### 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if



$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

## 5. Answer: d

**Explanation:**

$$(2x)^{2y} = 4e^{2x-2y}$$

$$2y \ln 2x = \ln 4 + 2x - 2y$$

$$y = \frac{x + \ln 2}{1 + \ln 2x}$$

$$y' = \frac{(1 + \ln 2x) - (x + \ln 2) \cdot \frac{1}{x}}{(1 + \ln 2x)^2}$$

$$y' = 1 + \ln 2x)^2 = \left[ \frac{x \ln 2x - \ln 2}{x} \right]$$

**Concepts:**

### 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

## 6. Answer: d

### Explanation:

$$f(x) = \sin(\sin x)$$

$$\Rightarrow f'(x) = \cos(\sin x) \cdot \cos x$$

$$\Rightarrow f''(x) = -\sin(\sin x) \cdot \cos^2(\sin x) \cdot (-\sin x)$$

$$= -\cos^2 x \cdot \sin(\sin x) - \sin x \cdot \cos(\sin x)$$

$$\text{Now } f''(x) + \tan x \cdot f'(x) + g(x) = 0$$

$$\Rightarrow g(x) = \cos^2 x \cdot \sin(\sin x) + \sin x \cdot \cos(\sin x) - \tan x \cdot \cos x \cdot \cos(\sin x)$$

$$\Rightarrow g(x) = \cos^2 x \cdot \sin(\sin x)$$

### Concepts:

#### 1. Continuity:

A function is said to be continuous at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

## 7. Answer: c

### Explanation:

Given:  $x = \sqrt{2^{\csc^{-1}t}}$  and  $y = \sqrt{2^{\sec^{-1}t}}$  ( $|t| \geq 1$ )

Now,  $\frac{dy}{dx} = \frac{dy/dt}{dx/dt}$

$$\begin{aligned}
 &= \frac{\frac{1}{2\sqrt{2^{\sec^{-1}t}}} 2^{\sec^{-1}t} \ln\left(\frac{1}{t\sqrt{t^2-1}}\right)}{-\frac{1}{2\sqrt{2^{\csc^{-1}t}}} 2^{\csc^{-1}t} \ln\left(\frac{1}{t\sqrt{t^2-1}}\right)} \\
 &= -\frac{\sqrt{2^{\sec^{-1}t}}}{\sqrt{2^{\csc^{-1}t}}} = \frac{-y}{x}
 \end{aligned}$$

### Concepts:

#### 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

---

## 8. Answer: d

**Explanation:**

$$\frac{dx}{dt} = 3 \sec^2 t$$

$$\frac{dy}{dt} = 3 \sec t \tan t$$

$$\frac{dy}{dx} = \frac{\tan t}{\sec t} = \sin t$$

$$\frac{d^2y}{dx^2} = \cos t \frac{dt}{dx}$$

$$= \frac{\cos t}{3 \sec^2 t} = \frac{\cos^3 t}{3} = \frac{1}{3 \cdot 2\sqrt{2}} = \frac{1}{6\sqrt{2}}$$

**Concepts:**

### 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

---

## 9. Answer: a

**Explanation:**

$$f : (-1, 1) \rightarrow R$$

$$f(x) = \max\{-|x|, -\sqrt{1-x^2}\}$$

Non-derivable at 3 points in  $(-1, 1)$

**Concepts:**

### 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function:** For any function to be continuous, it must meet the following conditions:

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .

## 10. Answer: c

### Explanation:

For continuity at  $x = 0$

$$\lim_{x \rightarrow \theta} \left\{ \frac{(e^x - 1)^2}{\sin\left(\frac{x}{k}\right) \cdot \ln\left(1 + \frac{x}{4}\right)} \right\} = 12$$

$$\Rightarrow \lim_{x \rightarrow 0} \left[ \frac{\left(\frac{e^x - 1}{x}\right)^2}{\frac{\sin\left(\frac{x}{k}\right)}{k\left(\frac{x}{k}\right)} \cdot \frac{\ln\left(1 + \frac{x}{4}\right)}{4 \cdot \left(\frac{x}{4}\right)}} \right]$$

$$\Rightarrow \left\{ \frac{(1)^2}{\left(\frac{1}{k}\right) \cdot \frac{1}{4}(1)} \right\} = 12$$

$$\Rightarrow 4k = 12 \Rightarrow k = 3$$

### Concepts:

#### 1. Continuity:

A function is said to be [continuous](#) at a point  $x = a$ , if

$$\lim_{x \rightarrow a}$$

$f(x)$  Exists, and

$$\lim_{x \rightarrow a}$$

$$f(x) = f(a)$$

It implies that if the left hand limit (L.H.L), right hand limit (R.H.L) and the value of the function at  $x=a$  exists and these parameters are equal to each other, then the function  $f$  is said to be continuous at  $x=a$ .

If the function is undefined or does not exist, then we say that the function is discontinuous.

**Conditions for continuity of a function: For any function to be continuous, it must meet the following conditions:**

- The function  $f(x)$  specified at  $x = a$ , is continuous only if  $f(a)$  belongs to real number.
- The limit of the function as  $x$  approaches  $a$ , exists.
- The limit of the function as  $x$  approaches  $a$ , must be equal to the function value at  $x = a$ .